

UNITED STATES PATENT APPLICATION

OF

URSULA SCHINDLER, KARL SCHÖNAFINGER, and HARTMUT STROBEL

FOR


**SULFUR SUBSTITUTED SULFONYLAMINOCARBOXYLIC ACID N-ARYLAMIDES,
THEIR PREPARATION, THEIR USE AND PHARMACEUTICAL PREPARATIONS
COMPRISING THEM**

Amend.
B¹

The present application claims priority under 35 U.S.C. § 119 to German patent applications No. 19830430.7 filed July 8, 1998, and No. 19903126.6 filed January 27, 1999. Both priority applications are entirely incorporated herein by reference.

5

The present invention relates to compounds of the formula I



15 wherein A¹, A², R¹, R², R³, X and n are as defined below, which are valuable pharmaceutically active compounds for the therapy and prophylaxis of diseases, for example for cardiovascular diseases such as hypertension, angina pectoris, cardiac insufficiency, thromboses or atherosclerosis. The compounds of the formula I are capable of modulating the body's production of cyclic guanosine monophosphate

20 ("cGMP") and are generally suitable for the therapy and prophylaxis of diseases which are associated with a disturbed cGMP balance. The invention furthermore relates to

processes for preparing compounds of the formula I, to their use for the therapy and prophylaxis of the abovementioned diseases and for preparing pharmaceuticals for this purpose, and to pharmaceutical preparations which comprise compounds of the formula I.

5

5
10
15
20

cGMP is an important intracellular messenger which triggers a multitude of different effects via the modulation of cGMP-dependent protein kinases, phosphodiesterases and ion channels. Examples are the relaxation of smooth muscles, the inhibition of thrombocyte activation and the inhibition of the proliferation of smooth-muscle cells and of leukocyte adhesion. cGMP is produced by particulate and soluble guanylate cyclases as a response to a number of extracellular and intracellular stimuli. In the case of the particulate guanylate cyclases, stimulation is essentially effected by peptidic messengers, such as the atrial natriuretic peptide or the cerebral natriuretic peptide. The soluble guanylate cyclases ("sGC"), which are cytosolic heterodimeric heme proteins, in contrast, are essentially regulated by a family of low-molecular-weight factors which are formed enzymatically. The most important stimulant is nitrogen monoxide ("NO") or a closely related species. The function of other factors such as carbon monoxide or the hydroxyl radical is still largely unclear. The binding of NO to the heme with formation of a penta-coordinate heme-nitrosyl complex is being discussed as activation mechanism of the activation by NO. The associated release of the histidine which is bound in the basal state to the iron converts the enzyme into the active conformation.

Active soluble guanylate cyclases are composed of an α and a β subunit each. Several subunit subtypes have been described which differ from one another with respect to sequence, tissue-specific distribution and expression in different development stages.

The subtypes α_1 and β_1 are mainly expressed in brain and lung, while β_2 is found in particular in liver and kidney. The subtype α_2 was shown to be present in human fetal brain. The subunits referred to as α_3 and β_3 were isolated from human brain and are homologous to α_1 and β_1 . More recent works indicate an α_{2i} subunit which contains an insert in the catalytic domain. All subunits show great homologies in the region of the catalytic domain. The enzymes presumably contain one heme per heterodimer, which is bound via β_1 -Cys-78 and/or β_1 -His-105 and is part of the regulatory center.

Under pathologic conditions, the formation of guanylate-cyclase-activating factors can be reduced, or their degradation may be promoted owing to the increased occurrence of free radicals. The resulting reduced activation of the sGC leads, via a weakening of the respective cGMP-mediated cellular response, for example to an increase of the blood pressure, to platelet activation or to increased cell proliferation and cell adhesion. As a consequence, formation of endothelial dysfunction, atherosclerosis, hypertension, stable or unstable angina pectoris, thromboses, myocardial infarction, strokes or erectile dysfunction results. Pharmacological stimulation of sGC offers a possibility to normalize

cGMP production and therefore makes possible the treatment and/or prevention of such disorders.

For the pharmacological stimulation of the sGC, use has hitherto almost exclusively

5 been made of compounds whose activity is based on an intermediate NO release, for example organic nitrates. The drawback of this treatment is the development of tolerance and a reduction of activity, and the higher dosage which is required because of this.

10 Various sGC stimulators which do not act via NO release were described by Vesely in a series of publications. However, the compounds, most of which are hormones, plant hormones, vitamins or natural compounds such as, for example, lizard poisons predominantly only have weak effects on the cGMP formation in cell lysates. D. L. Vesely, *Eur. J. Clin. Invest.*, vol. 15, 1985, p. 258; D. L. Vesely, *Biochem. Biophys. Res. Comm.*, vol. 88, 1979, p. 1244. A stimulation of heme-free guanylate cyclase by
15 protoporphyrin IX was demonstrated by Ignarro et al., *Adv. Pharmacol.*, vol. 26, 1994, p. 35. Pettibone et al., *Eur. J. Pharmacol.*, vol. 116, 1985 p. 307, described an antihypertensive action of diphenyliodonium hexafluorophosphate and attributed this to a stimulation of sGC. According to Yu et al., *Brit. J. Pharmacol.*, vol. 114, 1995, p. 1587,
20 isoliquiritigenin, which has a relaxing action on isolated rat aortas, also activates sGC. Ko et al., *Blood* vol. 84, 1994, p. 4226, Yu et al., *Biochem. J.* vol. 306, 1995, p. 787, and


Wu et al., *Brit. J. Pharmacol.* vol. 116, 1995, p. 1973, demonstrated a sGC-stimulating activity of 1-benzyl-3-(5-hydroxymethyl-2-furyl)indazole and demonstrated an antiproliferative and thrombocyte-inhibiting action. Pyrazoles and fused pyrazoles which exhibit a sGC-stimulating activity are described in European Patent Application No. 5 908,456 and German Patent Application No. 19,744,027.

A series of 2-sulfonylaminobenzoic acid N-arylamides, the N-aryl group of which carries a thio substituent, have been mentioned in the literature. These compounds in which the N-aryl group generally carries as further substituents groups which are readily oxidizable
10 such as, for example, two hydroxy groups being in para position with respect to one another and which in this case can be regarded as hydroquinone derivatives, are auxiliaries for the preparation of photographic materials (see, for example, Chemical Abstracts 119, 105757; 120, 41858; 123, 70224; or 126, 257007). If isolated structural elements are considered then the N-aryl group in these known compounds corresponds
15 to the group $R^1-S(O)_n-A^1$ in formula I in case A^1 denotes a 1,4-phenylene residue which in positions 2 and 5 carries hydroxy groups (or oxy substituents), and the number n is 0. British patent publication No. 876,526 (Chemical Abstracts 56, 15432e) discloses 3,5-dichloro-2-methylsulfonylaminobenzoic acid N-(5-chloro-2-(4-chlorophenylmercapto)-phenyl)-amide which can be used for the protection of wool against moths. Compounds
20 covered by British patent publication No. 876,526 correspond to compounds of the formula I if simultaneously the ring A^1 which comprises the carbon atoms which carry

the groups $C(=X)-NH-$ and $NH-SO_2R^2$, together with the residues R^3 , is a benzene ring which carries one to four halogen atoms from the series chlorine and bromine, R^2 is (C_1-C_4) -alkyl, X is oxygen and the group $R^1-S(O)_n-A^1-$ is a phenylmercaptophenyl- residue (= phenylthiophenyl-) which is substituted by halogen and/or trifluoromethyl and which can also be substituted by methyl or (C_1-C_4) -alkoxy, and the total number of halogen atoms and trifluoromethyl groups is greater than two. Pharmacological activities of these known 2-sulfonylaminobenzoic acid N-arylamides are not disclosed.

Surprisingly, it has now been found that the compounds of the present invention effect a strong activation of guanylate cyclase and are therefore suitable for the therapy and prophylaxis of disorders which are associated with a low cGMP level.

Thus, the present invention relates to compounds of the formula I:



wherein

A¹ is a divalent residue from the series phenylene, naphthylene and heteroarylene which can all be substituted by one or more identical or different substituents from the series halogen, (C₁-C₅)-alkyl, phenyl, tolyl, CF₃, NO₂, OH, -O-(C₁-C₅)-alkyl, -O-(C₂-C₄)-alkyl-O-(C₁-C₃)-alkyl, (C₁-C₂)-alkylenedioxy, NH₂, -NH-(C₁-C₃)-alkyl, -N((C₁-C₃)-alkyl)₂, -NH-CHO, -NH-CO-(C₁-C₅)-alkyl, -CN, -CO-NH₂, -CO-NH-(C₁-C₃)-alkyl, -CO-N((C₁-C₃)-alkyl)₂, -CO-OH, -CO-O-(C₁-C₅)-alkyl, heterocyclyl, CHO, -CO-(C₁-C₅)-alkyl, -S(O)_n-(C₁-C₄)-alkyl, -S(O)_n-phenyl and -S(O)_n-tolyl;

the ring A² which comprises the carbon atoms which carry the groups C(=X)-NH- and

NH-SO₂R² is a benzene ring, a naphthalene ring, a saturated or partially unsaturated 3-membered to 7-membered carbocycle, a saturated or partially unsaturated or aromatic

monocyclic 5-membered to 7-membered heterocycle which contains one or more ring heteroatoms from the series N, O and S, or a saturated or partially unsaturated or aromatic bicyclic 8-membered to 10-membered heterocycle which contains one or more ring heteroatoms from the series N, O and S;

5

R¹ is aryl, heterocyclyl or (C₁-C₁₈)-alkyl which can be substituted by one or more identical or different residues R⁴ or, if the number n in the group R¹-S(O)_n- is 2, R¹ can also be NR⁵R⁶ or, if the number n in the group R¹-S(O)_n- is 0, R¹ can also be -CN;

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995

R² is aryl, heterocyclyl, NR⁵R⁶ or (C₁-C₁₀)-alkyl which can be substituted by one or more identical or different residues R⁴;

R³ denotes one or more identical or different residues from the series hydrogen, halogen, CF₃, OH, -O-(C₁-C₇)-alkyl, -O-(C₂-C₄)-alkyl-O-(C₁-C₇)-alkyl, -O-aryl, (C₁-C₂)-alkylenedioxy, NO₂, -CN, NR⁷R⁸, -CO-NR⁷R⁸, -CO-OH, -CO-O-(C₁-C₅)-alkyl, heterocyclyl, -S(O)_n-(C₁-C₅)-alkyl and (C₁-C₅)-alkyl which can be substituted by one or more identical or different residues R⁴;

R⁴ is fluorine, OH, -O-(C₁-C₁₀)-alkyl, -O-(C₂-C₄)-alkyl-O-(C₁-C₇)-alkyl, -O-aryl, -CN, NR⁷R⁸, -CO-NH₂, -CO-NH-(C₁-C₃)-alkyl, -CO-N((C₁-C₃)-alkyl)₂, -CO-OH, -CO-O-(C₁-C₅)-alkyl, heterocyclyl or oxo;

R^5 is hydrogen, (C_1-C_{10}) -alkyl which can be substituted by one or more identical or different substituents R^4 and/or by aryl, or is aryl, heterocyclyl, $-CO-NR^7R^8$, $-CO$ -aryl or $-CO-(C_1-C_{10})$ -alkyl wherein the alkyl residue can be substituted by one or more identical or different residues R^4 ;

5

R^6 independently of R^5 has one of the meanings indicated for R^5 , or

R^5 and R^6 together with the nitrogen atom to which they are bonded form a 5-membered to 8-membered saturated or partially unsaturated ring which in addition to the nitrogen atom which carries the groups R^5 and R^6 can contain one or more further ring

10 heteroatoms from the series N, O and S and which can be substituted by one or more identical or different substituents from the series fluorine, (C_1-C_5) -alkyl, hydroxy- (C_1-C_3) -alkyl-, $-(C_1-C_3)$ -alkyl-O- (C_1-C_4) -alkyl, aryl, CF_3 , OH, $-O-(C_1-C_7)$ -alkyl, $-O$ -aryl, $-O-(C_2-C_4)$ -alkyl-O- (C_1-C_7) -alkyl, (C_2-C_3) -alkylenedioxy, NR^7R^8 , $-CN$, $-CO-NH_2$, $-CO-NH-(C_1-C_3)$ -alkyl, $-CO-N((C_1-C_3)$ -alkyl) $_2$, $-CO-OH$, $-CO-O-(C_1-C_5)$ -alkyl, CHO, $-CO-(C_1-C_5)$ -alkyl,

15 $-S(O)_n-(C_1-C_4)$ -alkyl, $-S(O)_n-NH_2$, $-S(O)_n-NH-(C_1-C_3)$ -alkyl, $-S(O)_n-N((C_1-C_3)$ -alkyl) $_2$, oxo, $-(CH_2)_m-NH_2$, $-(CH_2)_m-NH-(C_1-C_4)$ -alkyl and $-(CH_2)_m-N((C_1-C_4)$ -alkyl) $_2$ where in the substituent $-(CH_2)_m-N((C_1-C_4)$ -alkyl) $_2$ the two alkyl groups can be connected by a single bond and then together with the nitrogen atom carrying them form a 5-membered to 7-membered ring which besides that nitrogen atom and the carbon atoms can additionally

20 contain an oxygen atom, a sulfur atom or a group NR^5 as ring member;

R⁷ is hydrogen or (C₁-C₇)-alkyl which can be substituted by one or more identical or different substituents from the series OH, -O-(C₁-C₅)-alkyl, NH₂, -NH-(C₁-C₄)-alkyl and -N((C₁-C₄)-alkyl)₂ where in the substituent N((C₁-C₄)-alkyl)₂ the two alkyl groups can be connected by a single bond and then together with the nitrogen atom carrying them

5 form a 5-membered to 7-membered ring which besides that nitrogen atom and the carbon atoms can additionally contain an oxygen atom, a sulfur atom or a group NR⁵ as ring member;

R⁸ independently of R⁷ has one of the meanings of R⁷ or is -CO-(C₁-C₄)-alkyl;

10 "aryl" is phenyl, naphthyl or heteroaryl which can all be substituted by one or more identical or different substituent from the series halogen, (C₁-C₅)-alkyl, phenyl, tolyl, CF₃, -O-CF₃, NO₂, OH, -O-(C₁-C₅)-alkyl, -O-(C₂-C₄)-alkyl-O-(C₁-C₃)-alkyl, (C₁-C₂)-alkylenedioxy, NH₂, -NH-(C₁-C₃)-alkyl, -N((C₁-C₃)-alkyl)₂, -NH-CHO, -NH-CO-(C₁-C₅)-alkyl, -CN, -CO-NH₂, -CO-NH-(C₁-C₃)-alkyl, -CO-N((C₁-C₃)-alkyl)₂, -CO-OH, -CO-O-(C₁-C₅)-alkyl, heterocyclyl, CHO, -CO-(C₁-C₅)-alkyl, -S(O)_n-(C₁-C₄)-alkyl, -S(O)_n-phenyl and -S(O)_n-tolyl;

"heteroaryl" and "heteroarylene" are a residue of a monocyclic 5-membered or 6-membered aromatic heterocycle or of a bicyclic 8-membered to 10-membered aromatic

heterocycle each of which contains one or more ring heteroatoms from the series N, O and S;

"heterocyclyl" is a residue of a monocyclic or polycyclic 5-membered to 11-membered saturated or partially unsaturated heterocycle which contains one or more ring heteroatoms from the series N, O and S and which can be substituted by one or more identical or different substituents from the series fluorine, (C₁-C₅)-alkyl, OH, -O-(C₁-C₅)-alkyl, -O-(C₂-C₄)-alkyl-O-(C₁-C₃)-alkyl, NH₂, -NH-(C₁-C₃)-alkyl, -N((C₁-C₃)-alkyl)₂, -CN, -CO-NH₂, -CO-NH-(C₁-C₃)-alkyl, -CO-N((C₁-C₃)-alkyl)₂, -CO-OH and -CO-O-(C₁-C₅)-alkyl;

n is 0, 1 or 2;

m is 2, 3 or 4;

X is O or NH or X is a nitrogen atom which via a single bond is attached to a ring carbon atom in the group A¹ which ring carbon atom is directly adjacent to the carbon atom in A¹ carrying the group -NH-C(=X)- so that the group -NH-C(=X)- together with the carbon atoms in A¹ carrying it forms an anellated imidazole ring;

in all their stereoisomeric forms and mixtures thereof in all ratios, and their physiologically acceptable salts;

where, however, the compound of formula I is excluded wherein simultaneously the ring A^2 which comprises the carbon atoms which carry the groups $C(=X)-NH-$ and $NH-SO_2R^2$ is a benzene ring which is substituted in positions 3 and 5 by chlorine, R^2 is methyl, X is oxygen and $R^1-S(O)_n-A^1-$ is a 5-chloro-2-(4-chlorophenylmercapto)-phenyl residue.

5

If groups or substituents can occur several times in the compounds of formula I such as, for example R^3 , R^4 , R^5 , aryl, heterocyclyl, alkyl, or the numbers n and m , they can all independently of one another have the meanings indicated and can in each case be identical or different.

10

Alkyl residues can be straight-chain or branched. This also applies when they are part of other groups, for example in alkoxy groups, alkoxycarbonyl groups or amino groups, or when they are substituted. Examples of alkyl groups are methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl,

15 pentadecyl, hexadecyl, heptadecyl, octadecyl, the n -isomers of these residues, isopropyl, isobutyl, isopentyl, sec-butyl, tert-butyl, neopentyl, 3,3-dimethylbutyl. The term alkyl here also expressly includes unsaturated alkyl residues, i. e. alkyl residues which contain one or more double bonds and/or one or more triple bonds such as, for example, alkenyl residues and alkynyl residues. Of course, an unsaturated alkyl group
20 contains at least two carbon atoms. Specific alkyl groups whose number of carbon atoms can vary from 1 to a given upper limit, thus also comprise unsaturated alkyl

groups whose number of carbon atoms can vary from 2 to the given upper limit.

Examples of such residues are the vinyl residue, the 1-propenyl residue, the 2-propenyl residue (allyl residue), the 2-butenyl residue, the 2-methyl-2-propenyl residue, the 3-methyl-2-butenyl residue, the ethinyl residue, the 2-propinyl residue (propargyl residue),
5 the 2-butylnyl residue or the 3-butylnyl residue. Further, the term alkyl here also expressly includes alkyl residues in which by an internal ring closure within the alkyl group a cyclic system is formed, i. e. the term alkyl also includes saturated and partially unsaturated cycloalkyl residues and cycloalkyl-alkyl- residues (alkyl substituted by cycloalkyl). Of course, a monocyclic cycloalkyl group contains at least three carbon atoms.

Specific alkyl groups whose number of carbon atoms can vary from 1 to a given upper limit, thus also comprise monocyclic cycloalkyl groups whose number of carbon atoms can vary from 3 to the given upper limit, and appropriate cycloalkyl-alkyl- groups.

Examples of such cycloalkyl residues are cyclopropyl, cyclobutyl, cyclopentyl,

15 cyclohexyl, cycloheptyl and cyclooctyl which can all also be substituted by one or more identical or different (C₁-C₄)-alkyl residues, in particular by methyl. Examples of such substituted cycloalkyl residues are 4-methylcyclohexyl, 4-tert-butylcyclohexyl or 2,3-dimethylcyclopentyl. Furthermore, unless stated otherwise the term alkyl here also expressly includes unsubstituted alkyl residues as well as alkyl residues which are

20 substituted by one or more, for example one, two, three or four, identical or different aryl residues. The term alkyl thus here also expressly includes arylalkyl- residues such as,

for example, aryl-(C₁-C₄)-alkyl-, for example benzyl residues, phenylethyl residues or indanyl residues. In substituted alkyl residues, for example arylalkyl-, hydroxyalkyl- such as -(C₁-C₃)-alkyl-OH or alkoxyalkyl- such as -(C₁-C₃)-alkyl-O-(C₁-C₄)-alkyl, the substituents can be present in any desired position.

5

A saturated or partially unsaturated 3-membered to 7-membered carbocycle representing the ring A² can be derived from the monocyclic parent systems cyclopropane, cyclobutane, cyclopentane, cyclohexane or cycloheptane. If the carbocycle is unsaturated it can contain, for example, one double bond or, in the case of a 5-membered ring, 6-membered ring or 7-membered ring, also two double bonds which can be isolated or conjugated. Double bonds can be present in any positions with respect to the groups C(=X)-NH- and NH-SO₂-R², i.e., for example a double bond can also be present between the two ring carbon atoms which carry these two groups.

15 Unless stated otherwise, phenyl residues, naphthyl residues and heterocyclic residues, for example heteroaryl residues, can be unsubstituted or can carry one or more, for example one, two, three or four, identical or different substituents which can be in any desired positions. Unless stated otherwise, in these residues for example those substituents can be present which are indicated as substituents of an aryl group. A
20 preferred series of substituents that can be present in the residue aryl is formed by the substituents halogen, (C₁-C₅)-alkyl, phenyl, tolyl, CF₃, NO₂, OH, -O-(C₁-C₅)-alkyl, -O-(C₂-

C₄)-alkyl-O-(C₁-C₃)-alkyl, (C₁-C₂)-alkylenedioxy, NH₂, -NH-(C₁-C₃)-alkyl, -N((C₁-C₃)-alkyl)₂, -NH-CHO, -NH-CO-(C₁-C₅)-alkyl, -CN, -CO-NH₂, -CO-NH-(C₁-C₃)-alkyl, -CO-N((C₁-C₃)-alkyl)₂, -CO-OH, -CO-O-(C₁-C₅)-alkyl, heterocyclyl, CHO, -CO-(C₁-C₅)-alkyl, -S(O)_n-(C₁-C₄)-alkyl, -S(O)_n-phenyl and -S(O)_n-tolyl. If in compounds of the formula

5 I nitro groups are present as substituents, in total only up to two nitro groups can be present in the molecules. If phenyl residues, phenoxy residues, benzyl residues or benzyloxy residues are present as substituents in, for example, aryl residues like phenyl residues and/or in heterocyclic residues then in these substituents the benzene ring can also be unsubstituted or substituted by one or more, for example one, two, three or four, identical or different residues, for example by residues from the series (C₁-C₄)-alkyl, 10 halogen, hydroxy, (C₁-C₄)-alkoxy, trifluoromethyl, cyano, hydroxycarbonyl, ((C₁-C₄)-alkoxy)carbonyl, aminocarbonyl, nitro, amino, (C₁-C₄)-alkylamino, di-((C₁-C₄)-alkyl)amino and ((C₁-C₄)-alkyl)carbonylamino.

15 In monosubstituted phenyl residues the substituent can be in the 2-position, the 3-position or the 4-position, in disubstituted phenyl residues the substituents can be in 2,3-position, 2,4-position, 2,5-position, 2,6-position, 3,4-position or 3,5-position. In trisubstituted phenyl residues the substituents can be in 2,3,4-position, 2,3,5-position, 2,3,6-position, 2,4,5-position, 2,4,6-position or 3,4,5-position. Toly (i.e., methylphenyl) 20 can be 2-tolyl, 3-tolyl or 4-tolyl. Naphthyl can be 1-naphthyl or 2-naphthyl. In monosubstituted 1-naphthyl residues the substituent can be in the 2-position, the 3-

position, the 4-position, the 5-position, the 6-position, the 7-position or the 8-position, in monosubstituted 2-naphthyl residues in the 1-position, the 3-position, the 4-position, the 5-position, the 6-position, the 7-position or the 8-position.

- 5 The above explanations as well as the following explanations relating to monovalent residues correspondingly apply to the divalent residues phenylene, naphthylene and heteroarylene. The free bonds via which the divalent residues are attached to the adjacent groups can be present on any ring carbon atoms. In the case of a phenylene residue they can be in 1,2-position (ortho-phenylene), 1,3-position (meta-phenylene) or 1,4-position (para-phenylene). In the case of a naphthylene residue the free bonds can be in 1,2-position (= 1,2-naphthylene or 1,2-naphthalenediyl) or in 1,3-position, 1,4-position, 1,5-position, 1,6-position, 1,7-position, 1,8-position, 2,3-position, 2,6-position or 2,7-position. In the case of 5-membered ring aromatics containing one heteroatom such as, for example, thiophene or furan, the two free bonds can be in 2,3-position, 2,4-position, 2,5-position or 3,4-position. A divalent residue derived from pyridine can be a 2,3-, 2,4-, 2,5-, 2,6-, 3,4- or 3,5-pyridinediyl residue. In the case of unsymmetrical divalent residues the present invention includes all positional isomers, i.e., in the case of a 2,3-pyridinediyl residue, for example, it includes the compound in which the one adjacent group is present in the 2-position and the other adjacent group is present in the 3-position as well as the compound in which the one adjacent group is present in the 3-position and the other adjacent group is present in the 2-position.
- 15
- 20

Heteroaryl residues, heteroarylene residues, heterocyclyl residues, heterocycles representing the ring A² and rings which are formed by two groups bonded to a nitrogen atom together with this nitrogen atom are preferably derived from heterocycles which contain one, two, three or four identical or different ring heteroatoms, more preferably
5 from heterocycles which contain one, two or three, in particular one or two, identical or different heteroatoms. Unless stated otherwise, the heterocycles can be monocyclic or polycyclic, for example monocyclic, bicyclic or tricyclic. Preferably they are monocyclic or bicyclic. The rings preferably are 5-membered rings, 6-membered ring or 7-membered ring. Examples of monocyclic and bicyclic heterocyclic systems from which
10 residues occurring in the compounds of the formula I can be derived, are pyrrole, furan, thiophene, imidazole, pyrazole, 1,2,3-triazole, 1,2,4-triazole, 1,3-dioxole, 1,3-oxazole (= oxazole), 1,2-oxazole (= isoxazole), 1,3-thiazole (= thiazole), 1,2-thiazole (= isothiazole), tetrazole, pyridine, pyridazine, pyrimidine, pyrazine, pyran, thiopyran, 1,4-dioxine, 1,2-oxazine, 1,3-oxazine, 1,4-oxazine, 1,2-thiazine, 1,3-thiazine, 1,4-thiazine, 1,2,3-triazine,
15 1,2,4-triazine, 1,3,5-triazine, 1,2,4,5-tetrazine, azepine, 1,2-diazepine, 1,3-diazepine, 1,4-diazepine, 1,3-oxazepine, 1,3-thiazepine, indole, benzothiophene, benzofuran, benzothiazole, benzimidazole, quinoline, isoquinoline, cinnoline, quinazoline, quinoxaline, phthalazine, thienothiophenes, 1,8-naphthyridine and other naphthyridines, pteridin, or phenothiazine, each of them in saturated form (perhydro form) or in partially
20 unsaturated form (for example in the dihydro form or the tetrahydro form) or in maximally unsaturated form, insofar as the respective forms are known and stable.

Thus, the heterocycles which are suitable also include, for example, the saturated heterocycles pyrrolidine, piperidine, piperazine, morpholine and thiomorpholine. The degree of saturation of heterocyclic groups is indicated in their individual definitions. Unsaturated heterocycles can contain, for example, one, two or three double bonds within the ring system. 5-membered rings and 6-membered rings can in particular also be aromatic.

The residues derived from these heterocycles can be attached via any suitable carbon atom. Nitrogen heterocycles which can carry a hydrogen atom or a substituent on a ring nitrogen atom, for example pyrrole, imidazole, pyrrolidine, morpholine, piperazine etc., can also be attached via a ring nitrogen atom, in particular if the heterocyclic residue in question is bonded to a carbon atom. For example, a thienyl residue can be present as 2-thienyl residue or 3-thienyl residue, a furyl residue as 2-furyl residue or 3-furyl residue, a pyridyl residue as 2-pyridyl residue, 3-pyridyl residue or 4-pyridyl residue, a piperidinyl residue as 1-piperidinyl residue (i.e., piperidino residue), 2-piperidinyl residue, 3-piperidinyl residue or 4-piperidinyl residue, a (thio)morpholinyl residue as 2-(thio)morpholinyl residue, 3-(thio)morpholinyl residue or 4-(thio)morpholinyl residue (i.e., thiomorpholino residue). A residue derived from 1,3-thiazole or imidazole which is attached via a carbon atom can be attached via the 2-position, the 4-position or the 5-position.

Unless stated otherwise the heterocyclic groups can be unsubstituted or can carry one or more, for example one, two, three, or four, identical or different substituents.

Substituents in heterocycles can be present in any desired positions, for example in a 2-thienyl residue or 2-furyl residue in the 3-position and/or in the 4-position and/or in the 5-

5 position, in a 3-thienyl residue or 3-furyl residue in the 2-position and/or in the 4-position and/or in the 5-position, in a 2-pyridyl residue in the 3-position and/or in the 4-position

and/or in the 5-position and/or in the 6-position, in a 3-pyridyl residue in the 2-position and/or in the 4-position and/or in the 5-position and/or in the 6-position, in a 4-pyridyl

10 residue in the 2-position and/or in the 3-position and/or in the 5-position and/or in the 6-position. Unless stated otherwise, for example those substituents can be present as

substituents in heterocyclic groups which are indicated in the definition of the group aryl, and in the case of saturated and partially unsaturated heterocycles as further

substituents also the oxo group and the thioxo group can be present. Substituents on a heterocycle as well as substituents on a carbocycle can also form a ring, i.e., to a ring

15 system further rings can be condensed (or anellated) so that, for example, also

cyclopenta-condensed, cyclohexa-condensed or benzo-condensed rings can be

present. Suitable substituents on a substitutable ring nitrogen atom of a heterocycle are

in particular, for example, unsubstituted (C₁-C₅)-alkyl residues and aryl-substituted alkyl residues, aryl residues, acyl residues such as -CO-(C₁-C₅)-alkyl, or sulfonyl residues

20 such as -SO₂-(C₁-C₅)-alkyl. Suitable nitrogen heterocycles can also be present as N-



oxides or as quaternary salts containing a counterion which is derived from a





physiologically acceptable acid. Pyridyl residues, for example, can be present as pyridine-N-oxides.



"Halogen" is fluorine, chlorine, bromine or iodine, and preferably fluorine or chlorine.

5

Without limiting the present invention, in the formulae la, lb, lc, ld, le, lf, lg and lh examples of groups of compounds of the invention are shown in which A² in the formula I has specific denotations. A¹, R¹, R², R³, X and n in the formulae la, lb, lc, ld, le, lf, lg and lh are defined as above for the formula I, and the number k in the formula lb is 1, 2, 3, 4 or 5, in particular 3 or 4.








10 On the benzene ring depicted in formula Ia which carries the groups C(=X)-NH- and
 -NHSO₂R², four positions are present which can carry a residue R³. The compounds of
 formula Ia can thus carry four residues R³ which, independently of one another, can all
 be hydrogen or can have a meaning different from hydrogen, i.e., in the compounds of
 formula Ia the benzene ring depicted in formula Ia can be unsubstituted or can carry
 15 one, two, three or four identical or different substituents from the series halogen, CF₃,
 OH, -O-(C₁-C₇)-alkyl, -O-(C₂-C₄)-alkyl-O-(C₁-C₇)-alkyl, -O-aryl, (C₁-C₂)-alkylenedioxy,
 NO₂, -CN, NR⁷R⁸, -CO-NR⁷R⁸, -CO-OH, -CO-O-(C₁-C₅)-alkyl, heterocyclyl, -S(O)_n-(C₁-
 C₅)-alkyl and (C₁-C₅)-alkyl which can be substituted by one or more identical or different
 residues R⁴. These explanations accordingly also apply to the compounds of formulae
 20 Ib to Ih.

The compounds of the formula I wherein X is a nitrogen atom which via a single bond is attached to a ring carbon atom in the group A¹ which ring carbon atom is directly adjacent to the carbon atom in A¹ carrying the group -NH-C(=X)- so that the group -NH-C(=X)- together with the carbon atoms in A¹ carrying it forms an anellated

5 imidazole ring, are represented by the formula li.



15 A², R¹, R², R³ and n in the formula li are defined as above for the formula I. The ring A³ which has resulted from the group A¹ by the formation of a bond to the nitrogen atom representing X and which ring comprises the two carbon atoms depicted in formula li carrying the nitrogen atoms of the anellated imidazole ring, is a benzene ring, a naphthalene ring or a heteroaromatic ring, where to these rings the above explanations

20 relating to A¹ correspondingly apply.

The present invention includes all stereoisomeric forms of the compounds of the formula

I. Centers of asymmetry that are present in the compounds of formula I can all independently of one another have S configuration or R configuration. The invention includes all possible enantiomers and diastereomers and mixtures of two or more

5 stereoisomers, for example mixtures of enantiomers and/or diastereomers, in all ratios.

Thus, enantiomers are a subject of the invention in enantiomerically pure form, both as levorotatory and as dextrorotatory antipodes, in the form of racemates and in the form of

mixtures of the two enantiomers in all ratios. In the case of a cis/trans isomerism the invention includes both the cis form and the trans form as well as mixtures of these

10 forms in all ratios. The preparation of individual stereoisomers can be carried out, if desired, by separation of a mixture by customary methods, for example by

chromatography or crystallization, by the use of stereochemically uniform starting

materials for the synthesis or by stereoselective synthesis. Optionally a derivatization

can be carried out before a separation of stereoisomers. The separation of a mixture of

15 stereoisomers can be carried out at the stage of the compounds of the formula I or at

the stage of an intermediate during the synthesis. The present invention also includes

all tautomeric forms of the compounds of formula I.

If the compounds of the formula I contain one or more acidic or basic groups the

20 invention also includes the corresponding physiologically or toxicologically acceptable

salts, in particular the pharmaceutically utilizable salts. Thus, the compounds of the

formula I which contain acidic groups can be present on these groups and can be used according to the invention, for example, as alkali metal salts, alkaline earth metal salts or as ammonium salts. Examples of such salts are sodium salts, potassium salts, calcium salts, magnesium salts or salts with ammonia or organic amines such as, for example, ethylamine, ethanolamine, triethanolamine or amino acids. Compounds of the formula I which contain one or more basic groups, i.e. groups which can be protonated, can be present and can be used according to the invention in the form of their acid addition salts with inorganic or organic acids, for example as salts with hydrogen chloride, hydrogen bromide, phosphoric acid, sulfuric acid, nitric acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acids, oxalic acid, acetic acid, tartaric acid, lactic acid, salicylic acid, benzoic acid, formic acid, propionic acid, pivalic acid, diethylacetic acid, malonic acid, succinic acid, pimelic acid, fumaric acid, maleic acid, malic acid, sulfaminic acid, phenylpropionic acid, gluconic acid, ascorbic acid, isonicotinic acid, citric acid, adipic acid, etc. If the compounds of the formula I simultaneously contain acidic and basic groups in the molecule the invention also includes, in addition to the salt forms mentioned, inner salts or betaines (zwitterions). Salts can be obtained from the compounds of the formula I by customary methods which are known to the person skilled in the art, for example by combination with an organic or inorganic acid or base in a solvent or dispersant, or by anion exchange or cation exchange from other salts. The present invention also includes all salts of the compounds of the formula I which, owing to low physiological compatibility, are not

directly suitable for use in pharmaceuticals but which can be used, for example, as intermediates for chemical reactions or for the preparation of physiologically acceptable salts.

- 5 The present invention furthermore includes all solvates of compounds of the formula I, for example hydrates or adducts with alcohols, and also derivatives of the compounds of the formula I such as, for example esters, amides, prodrugs and active metabolites.

A¹ preferably is a phenylene residue or a 5-membered or 6-membered heteroarylene residue, more preferably phenylene, where all these residues can be unsubstituted or can be substituted as indicated. If the group A¹ is substituted, i. e. if it carries one or more further substituents in addition to the group R¹-S(O)_n, it is preferably substituted by one or two identical or different of the substituents indicated above. Preferably a phenylene residue representing A¹ is unsubstituted, i. e. besides the groups R¹-S(O)_n and C(=X)-NH it carries four hydrogen atoms. The group R¹-S(O)_n is preferably attached to a carbon atom in A¹ which is not directly adjacent to the carbon atom which carries the group C(=X)-NH. If A¹ is phenylene the group R¹-S(O)_n is particularly preferably located in the meta position or in the para position, more particularly preferably in the para position, with respect to the carbon atom which carries the group C(=X)-NH.

The ring A² which comprises the two carbon atoms which carry the groups R²-SO₂-NH and C(=X)-NH- preferably is an aromatic ring, more preferably a benzene ring or a thiophene ring, particularly preferably a benzene ring, where all these rings can be unsubstituted or substituted by one or more residues R³ which are different from
5 hydrogen.

R¹ preferably is (C₁-C₇)-alkyl, NR⁵R⁶ or aryl, more preferably NR⁵R⁶, phenyl or 5-membered or 6-membered heteroaryl, particularly preferably NR⁵R⁶, where all these residues can be unsubstituted or substituted as indicated and where, as stated above, R¹ can be NR⁵R⁶ if the number n in the group R¹-S(O)_n- is 2.

R² preferably is aryl, more preferably phenyl or 5-membered or 6-membered heteroaryl, particularly preferably phenyl or a residue of a monocyclic 5-membered or 6-membered aromatic heterocycle which contains one or two identical or different heteroatoms from
15 the series N, O and S such as, for example, phenyl, thienyl, pyrazolyl, imidazolyl, isoxazolyl, thiazolyl, pyridyl etc., in particular phenyl or 2-thienyl, where all these residues can be unsubstituted or substituted as indicated. Preferably an aryl residue representing R² is substituted. If an aryl residue representing R² is substituted it is preferably substituted by one, two or three, in particular by one or two, identical or
20 different substituents. Substituents in an aryl residue representing R² preferably are substituents from the series halogen, CF₃, -O-CF₃, NO₂, -CN, (C₁-C₄)-alkyl and -O-(C₁-

C₄)-alkyl, more preferably substituents from the series F, Cl, Br, CF₃, -O-CF₃, NO₂, -CN, CH₃ and -OCH₃. Especially preferably a substituted aryl residue representing R² is substituted by Cl, for example by one or two, in particular one, chlorine atoms.

- 5 The rings representing A² can be unsubstituted or substituted as indicated. When they are unsubstituted they only carry residues R³ which are hydrogen. When they are substituted they carry one or more residues R³ which are different from hydrogen. Those substituent positions which do not carry a residue R³ which is different from hydrogen, carry hydrogen atoms. If the ring A² carries one or more residues R³ which are different from hydrogen it preferably carries one or two such residues R³, in particular one such residue R³. Residues R³ which are different from hydrogen are preferably located in positions of the ring A² which are not directly adjacent to the groups C(=X)-NH and R²-SO₂-NH. If A² is a saturated or partially unsaturated carbocycle, residues R³ which are different from hydrogen preferably are (C₁-C₄)-alkyl, in particular methyl. If A² is an aromatic ring, in particular if A² is a benzene ring, residues R³ which are different from hydrogen preferably are (C₁-C₃)-alkyl, halogen, (C₁-C₃)-alkoxy or CF₃, more preferably methyl, chlorine or methoxy. If A² is an aromatic ring, in particular a benzene ring, it is very particularly preferred if the ring carries one chlorine atom or two methoxy groups as substituents, i. e., if one residue R³ is present which is chlorine or if two residues R³ are present which are methoxy, and the other positions on the benzene ring carry hydrogen atoms. If A² is a benzene ring residues R³

which are different from hydrogen are preferably located in positions 4 and/or 5 (with respect to the group $C(=X)-NH$ in the 1-position and the group R^2-SO_2-NH in the 2-position).

5 If a group is substituted by one or more residues R^4 it is preferably substituted by one, two or three, in particular one or two, identical or different residues R^4 . R^4 preferably is hydroxy, (C_1-C_4) -alkyloxy, di- $((C_1-C_4)$ -alkyl)amino or heteroaryl.

10 R^5 and R^6 preferably are independently of one another hydrogen, (C_1-C_9) -alkyl, (C_1-C_4) -alkyl-O- (C_1-C_3) -alkyl- or 5-membered or 6-membered aryl or together with the nitrogen atom carrying R^5 and R^6 form a 5-membered to 7-membered heterocycle which in addition to the nitrogen atom carrying the groups R^5 and R^6 can contain one further ring heteroatom from the series N, O and S and which can be substituted by one or more, for example one, two, three or four, identical or different residues from the series $(C_1-$
15 $C_3)$ -alkyl, hydroxy- (C_1-C_3) -alkyl-, 5-membered or 6-membered aryl, carbamoyl, hydroxy and oxo. It is particularly preferred if R^5 and R^6 together with the nitrogen atom carrying these residues form a 5-membered, 6-membered or 7-membered heterocycle which in addition to the nitrogen atom carrying the groups R^5 and R^6 can contain one further ring heteroatom from the series N, O and S and which can be substituted by one or more,
20 for example one, two, three or four, identical or different residues from the series (C_1-C_3) -alkyl, hydroxy- (C_1-C_3) -alkyl-, 5-membered or 6-membered aryl, carbamoyl, hydroxy

and oxo, in particular (C₁-C₃)-alkyl such as, for example, methyl. Preferably a heterocycle which is formed from the groups R⁵ and R⁶ together with the nitrogen atom carrying these residues is saturated.

- 5 An especially preferable heterocycle which is formed by R⁵ and R⁶ together with the nitrogen atom carrying these residues, is derived from morpholine, thiomorpholine, 1,1-dioxo-thiomorpholine, 1-oxo-thiomorpholine, from dialkylmorpholines such as dimethylmorpholines, from 2,6-dimethylmorpholine, cis-2,6-dimethylmorpholine, 3,5-dimethylmorpholine, cis-3,5-dimethylmorpholine, 1-(pyrimidin-2-yl)-piperazine,
10 piperidine-4-carboxamide, 1-(2-hydroxyethyl)-piperazine, 1-methylpiperazine, 1-ethylpiperazine, from 1-aryl piperazines, from ethyl piperazine-1-carboxylate, piperidine, 2-methylpiperidine, 2,6-dimethylpiperidine, cis-2,6-dimethylpiperidine, 3,5-dimethylpiperidine, cis-3,5-dimethylpiperidine, 4-hydroxypiperidine, from 4-oxopiperidine or a ketal thereof like 1,4-dioxo-8-aza-spiro[4.5]decan, from tetrahydropyridine,
15 tetrahydropyrimidine, 1-methylhomopiperazine, thiazolidine, pyrroline, pyrrolidine, 3-hydroxypyrrolidine, 1,2,3,4-tetrahydroisoquinoline or 2,3-dihydro-1H-isoindole, where these rings are attached via the ring nitrogen atom or, in the case of piperazine derivatives, via the unsubstituted ring nitrogen atom. A more especially preferable heterocycle which is formed by R⁵ and R⁶ together with the nitrogen atom carrying these
20 residues, is derived from morpholine, thiomorpholine, 1,1-dioxo-thiomorpholine, 1-oxo-thiomorpholine, 2,6-dimethylmorpholine, cis-2,6-dimethylmorpholine, 3,5-

dimethylmorpholine, cis-3,5-dimethylmorpholine, 1-(pyrimidin-2-yl)-piperazine, piperidine-4-carboxamide, 1,2,3,4-tetrahydroisoquinoline or 2,3-dihydro-1H-isoindole, moreover preferably from morpholine, 2,6-dimethylmorpholine or cis-2,6-dimethylmorpholine, in particular from morpholine or cis-2,6-dimethylmorpholine, where
5 these rings are attached via the ring nitrogen atom or, in the case of the piperazine derivative, via the unsubstituted ring nitrogen atom.

R⁷ preferably is hydrogen, (C₁-C₃)-alkyl, ((C₁-C₄)-alkyl)₂N-(C₁-C₃)-alkyl- or (C₁-C₄)-alkyl-O-(C₁-C₃)-alkyl-.

R⁸ preferably is hydrogen, (C₁-C₃)-alkyl or acetyl.

Aryl preferably is phenyl or heteroaryl, in particular phenyl or 5-membered or 6-membered heteroaryl. Unless stated otherwise, preferred substituents on aryl residues
15 are halogen, CF₃, (C₁-C₃)-alkyl, cyano, nitro and (C₁-C₃)-alkyloxy, more preferred substituents are CF₃, chlorine, methyl and methoxy.

Heteroaryl and heteroarylene preferably are a residue of a monocyclic 5-membered or 6-membered aromatic heterocycle, in particular a residue derived from the
20 heteroaromatics thiophene, pyrazole, thiazole, oxazole, isoxazole, pyridine, pyrimidine, pyridazine and tetrazole.

Heterocyclyl preferably is a residue derived from a saturated heterocycle, more preferably a residue of a monocyclic 5-membered or 6-membered saturated heterocycle, in particular a residue which is derived from pyrrolidine, piperidine, from N-alkylpiperazines, from morpholine, from dialkylmorpholines, from thiomorpholine or tetrahydrofuran. In addition, the above explanations on preferred heterocycles which are formed by the residues R^5 and R^6 together with the nitrogen atom carrying these residues correspondingly apply to heterocyclyl residues which are attached via a ring nitrogen atom.

If a group $S(O)_n$ is bonded to a nitrogen atom the number n therein preferably is 1 or 2, more preferably 2. The number n in the group $R^1-S(O)_n$ preferably is 0 or 2, particularly preferably 2.

X preferably is O or a nitrogen atom which via a single bond is attached to a ring carbon atom in the group A^1 which ring carbon atom is directly adjacent to the carbon atom in A^1 carrying the group $-NH-C(=X)-$ so that the group $-NH-C(=X)-$ together with the carbon atoms in A^1 carrying it forms an anellated imidazole ring. Particularly preferably X is O.

Preferred compounds of the formula I are those compounds in which one or more of the residues contained therein have preferred meanings, all combinations of preferred

substituent definitions being a subject of the present invention. Also with respect to all preferred compounds of the formula I the present invention includes all stereoisomeric forms and mixtures thereof in all ratios, and their physiologically acceptable salts.

Groups of preferred compounds are also formed, for example, by the compounds of the

5 formulae Ia, Ib, Ic, Id, Ie, If, Ig and Ih in which one or more residues have preferred meanings, in all their stereoisomeric forms and mixtures thereof in all ratios, and their physiologically acceptable salts.

A group of particularly preferred compounds is formed, for example, by compounds of the formula I in which

A¹ is phenylene or heteroarylene where these residues can be unsubstituted or substituted by one or more identical or different substituents from the series halogen, (C₁-C₄)-alkyl, CF₃, -O-(C₁-C₄)-alkyl and -CN;

15 the ring A² which comprises the two carbon atoms which carry the groups R²-SO₂-NH and C(=X)-NH- is an aromatic ring;

R¹ is (C₁-C₇)-alkyl which can be substituted by one or more identical or different residues R⁴, or is aryl, or if the number n in the group R¹-S(O)_n- is 2 also is NR⁵R⁶;

20 R² is aryl;

R^3 denotes one or more identical or different residues from the series hydrogen, halogen, CF_3 , OH, $-O-(C_1-C_4)\text{-alkyl}$, $-O-(C_2-C_4)\text{-alkyl-O-(C}_1\text{-C}_4\text{)-alkyl}$, $-O\text{-aryl}$, NO_2 , $-CN$, NR^7R^8 , $-CO\text{-}NR^7R^8$, $-CO\text{-OH}$, $-CO\text{-O-(C}_1\text{-C}_4\text{)-alkyl}$, heterocyclyl, $-S(O)_n\text{-(C}_1\text{-C}_4\text{)-alkyl}$ and $(C_1-C_4)\text{-alkyl}$ which can be substituted by one or more identical or different residues R^4 ;

5

R^4 is fluorine, OH, $-O-(C_1-C_{10})\text{-alkyl}$, $-O-(C_2-C_4)\text{-alkyl-O-(C}_1\text{-C}_7\text{)-alkyl}$, $-O\text{-aryl}$, $-CN$, NR^7R^8 , $-CO\text{-NH}_2$, $-CO\text{-NH-(C}_1\text{-C}_3\text{)-alkyl}$, $-CO\text{-N}((C_1-C_3)\text{-alkyl})_2$, $-CO\text{-OH}$, $-CO\text{-O-(C}_1\text{-C}_4\text{)-alkyl}$, heterocyclyl or oxo;

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
214

"aryl" is phenyl or heteroaryl which can all be substituted by one or more identical or different substituents from the series halogen, (C₁-C₄)-alkyl, phenyl, CF₃, NO₂, -O-(C₁-C₄)-alkyl, (C₁-C₂)-alkylenedioxy, NH₂, -NH-CO-(C₁-C₄)-alkyl, -CN, -CO-NH₂, -CO-OH and -CO-O-(C₁-C₄)-alkyl;

5

heteroaryl and heteroarylene are a residue of a monocyclic 5-membered or 6-membered aromatic heterocycle which contains one or more identical or different ring heteroatoms from the series N, O and S;

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185

A group of more particularly preferred compounds is formed, for example, by compounds of the formula I in which:

A¹ is phenylene which is unsubstituted or substituted by one or more identical or different substituents from the series halogen, (C₁-C₄)-alkyl, CF₃, -O-(C₁-C₄)-alkyl and
5 -CN;

the ring A² which comprises the two carbon atoms which carry the groups R²-SO₂-NH and C(=X)-NH- is a benzene ring;

10 R¹ is NR⁵R⁶;

R² is aryl;

R³ denotes one or more identical or different residues from the series hydrogen,

15 halogen, CF₃, -O-(C₁-C₄)-alkyl, -CN and (C₁-C₄)-alkyl;

R⁵ and R⁶ together with the nitrogen atom carrying R⁵ and R⁶ form a 5-membered or 6-membered saturated heterocycle which in addition to the nitrogen atom carrying the groups R⁵ and R⁶ can contain one further ring heteroatom from the series N, O and S

20 and which can be substituted by one or more identical or different residues from the series (C₁-C₃)-alkyl, hydroxy-(C₁-C₃)-alkyl-, aryl, carbamoyl, hydroxy and oxo;

aryl is phenyl or 5-membered or 6-membered heteroaryl containing one or more identical or different ring heteroatoms from the series N, O and S which residues can all be substituted by one or more identical or different substituents from the series halogen, (C₁-C₄)-alkyl, CF₃, NO₂, -O-(C₁-C₄)-alkyl, -NH-CO-(C₁-C₄)-alkyl and -CN;

5

n is 2;

X is oxygen;

10 in all their stereoisomeric forms and mixtures thereof in all ratios, and their physiologically acceptable salts.

A group of especially preferred compounds is formed, for example, by compounds of the formula I in which:

15 A¹ is an unsubstituted divalent phenylene residue;

the ring A² which comprises the two carbon atoms which carry the groups R²-SO₂-NH and C(=X)-NH- is a benzene ring;

20 R¹ is NR⁵R⁶;

R² is aryl;

R³ denotes one or more identical or different residues from the series hydrogen, halogen, -O-(C₁-C₄)-alkyl and (C₁-C₄)-alkyl;

5

R⁵ and R⁶ together with the nitrogen atom carrying R⁵ and R⁶ form a saturated 6-membered heterocycle which in addition to the nitrogen atom carrying the groups R⁵ and R⁶ can contain one further ring heteroatom from the series N, O and S and which can be substituted by one or more identical or different residues from the series (C₁-C₃)-alkyl, aryl, oxo and carbamoyl;

aryl is phenyl or 5-membered or 6-membered heteroaryl containing one or more identical or different ring heteroatoms from the series N, O and S which residues can all be substituted by one or more identical or different substituents from the series halogen, (C₁-C₄)-alkyl, CF₃ and -O-(C₁-C₄)-alkyl;

n is 2;

X is oxygen;

20

in all their stereoisomeric forms and mixtures thereof in all ratios, and their physiologically acceptable salts.

A group of even more preferred compounds is formed, for example, by compounds of the formula I in which:

A¹ is an unsubstituted divalent 1,4-phenylene residue;

the ring A² which comprises the two carbon atoms which carry the groups R²-SO₂-NH and C(=X)-NH-, together with the residues R³, is a benzene ring which carries one or two substituents from the series chlorine and methoxy;

R¹ is NR⁵R⁶;

R² is phenyl or thienyl which residues are all substituted by one or two chlorine atoms;

R⁵ and R⁶ together with the nitrogen atom carrying R⁵ and R⁶ form a saturated 6-membered heterocycle which in addition to the nitrogen atom carrying the groups R⁵ and R⁶ can contain one further ring heteroatom from the series O and S and which is unsubstituted or substituted by one or two methyl residues;

n is 2;

X is oxygen;

in all their stereoisomeric forms and mixtures thereof in all ratios, and their physiologically acceptable salts.

5

The presents invention also relates to processes for the preparation of the compounds of the formula I which are described in the following and by which the compounds of the invention are obtainable. According to Scheme 1 compounds of the invention can be obtained, for example, by first reacting an aminocarboxylic acid of the formula II with a sulfonyl chloride of the formula R^2-SO_2-Cl or a sulfonic acid anhydride in the presence of a base in a solvent like water, pyridine or an ether. Suitable bases are inorganic bases like sodium carbonate or organic bases like, for example, pyridine or triethylamine. The sulfonylaminocarboxylic acid of the formula III that is obtained can then be activated, for example by reaction with a chlorinating agent like phosphorus pentachloride, phosphorus oxychloride or thionyl chloride in an inert solvent, to give an acid chloride of the formula IV which is then reacted with an arylamine.

15

The activation of the carboxylic acid group in the compounds of the formula III can, however, also be carried out in a different manner, for example by one of the numerous methods for the formation of amide bonds in peptide chemistry which are well-known to

20


the skilled person, for example by conversion into a mixed anhydride or an activated ester or by using a carbodiimide like dicyclohexylcarbodiimide.

The reaction of the activated sulfonylaminocarboxylic acid with an arylamine is favorably
 5 carried out in an inert solvent such as, for example, pyridine, tetrahydrofuran or toluene
 in the absence or in the presence of an inert auxiliary base like, for example, a tertiary
 amine or pyridine. If the arylamine that is employed in the reaction with the activated
 carboxylic acid already contains the desired substituent $R^1-S(O)_n$ then the reaction
 directly provides the final compound of the formula I. Compounds of the formula I in
 10 which the number n in the group $R^1-S(O)_n$ is 1 or 2 can also be obtained by reacting an
 activated carboxylic acid with a mercapto substituted arylamine of the formula R^1-S-A^1-
 NH_2 and then oxidizing the mercapto group in the compound of the formula V under
 standard conditions, for example with a peroxide like hydrogen peroxide or a peracid
 like 3-chloroperbenzoic acid or monoperoxyphthalic acid in a solvent like, for example,
 15 methylene chloride or acetone. The activated carboxylic acids can also first be reacted
 with arylamines of the formula A^1-NH_2 . The resulting reaction product of the formula VI
 can then be chlorosulfonated under standard conditions and the chlorosulfonyl group
 can then be converted under standard conditions into the group R^1-SO_2 , for example by
 reaction with suitable amines in substance or in a solvent like N-methylpyrrolidone,
 20 dimethylformamide, toluene or an ether, optionally in the presence of an auxiliary base.
 In a similar manner the activated carboxylic acids can be reacted with

fluorosulfonylarylamines of the formula $F-SO_2-A^1-NH_2$ and the fluorosulfonyl intermediates of the formula VII that are obtained can be converted under standard conditions into the compounds of the formula I according to the invention.


[illegible]

(Scheme 1)




The compounds of the formula I according to the invention can furthermore be obtained by reacting the activated sulfonylaminocarboxylic acids, for example the acid chlorides of the formula IV shown in Scheme 1, with a mercaptoarylamine of the formula H_2N-A^1-SH which is unsubstituted on the sulfur atom. In a nucleophilic substitution reaction the product of the formula VIII that is obtained can subsequently be alkylated or arylated on the sulfur atom with an alkyl halogenide or an aryl halogenide or another reactive compound under standard conditions and, if desired, be oxidized on the sulfur to give the sulfoxide or the sulfone as explained above with respect to the compounds of the formula V (see Scheme 2).

(Scheme 2)



Compounds of the formula I can also be obtained, for example, by first activating a suitably substituted nitrocarboxylic acid of the formula IX, for example by converting it into the respective acid chloride of the formula X or by another procedure, and then reacting it with a substituted arylamine of the formula $R^1-S(O)_n-A^1-NH_2$ analogously to
5 the procedures described above (see Scheme 3). Here, too, as arylamines fluorosulfonylarylamines of the formula $F-SO_2-A^1-NH_2$ can be employed, and in the N-(fluorosulfonylaryl)-carboxamides of the formula XI that are obtained the fluorosulfonyl group can be converted under standard conditions into a group R^1-SO_2 according to the invention, for example with an amine of the formula HNR^5R^6 .

(Scheme 3)



Before the nitro group is reduced to an amino group, in the nitro intermediates of the formula XII, use can be made of the activating effect of the nitro group on the ring A² and a suitable residue R³, for example a halogen atom, can be replaced with a different residue R³ by reaction with a nucleophile, for example an amine. The reduction of the

5 nitro group to give an amino group can, for example, be carried out by catalytic hydrogenation in the presence of a noble metal catalyst or, preferably, in the presence of Raney nickel in a solvent like ethanol, glacial acetic acid or an ethanolic solution of hydrogen chloride, or it can be carried out by reduction with a base metal like zinc, tin or iron in the presence of an acid. The reduction can also be carried out, for example, with

10 tin(II) chloride or by reaction with sodium dithionite, favorably in a mixture of methanol, tetrahydrofuran and water as solvent. The sulfonylation of the amino group in the reduction product of the formula XIII with an activated sulfonic acid derivative can be carried out analogously to the reactions described above, for example with a sulfonic acid chloride in pyridine, and finally gives the compound of the formula I.


15

The compounds of the formula I in which X is a nitrogen atom which via a single bond is attached to a ring carbon atom in the group A¹ which ring carbon atom is directly adjacent to the carbon atom in A¹ carrying the group -NH-C(=X)-, i. e. the benzimidazole derivatives of the formula II, can for example be obtained by reacting an

20 activated sulfonylaminocarboxylic acid derivative obtained as described above according to Scheme 1, for example a carboxylic acid chloride of the formula IV, (or

also, in analogy to Scheme 3, a nitrocarboxylic acid derivative) with a 1,2-diaminoarene in the presence of a dehydrating agent such as, for example, thionyl chloride or phosphorus pentachloride (see Scheme 4). The reaction is usually carried out in an inert solvent, for example in a hydrocarbon like toluene or xylene. The 1,2-diaminoarene can
5 already contain the final group $R^1-S(O)_n$ or a precursor group thereof, for example the group R^1-S . Subsequent steps, for example reactions on the sulfur atom, can then be carried out as explained above. Just so, unsubstituted 1,2-diaminoarenes can be employed and the resulting products of the formula XIV can be chlorosulfonated, for example with chlorosulfuric acid, and the sulfonyl chlorides that are obtained can be
10 converted into the final compounds containing the group R^1-SO_2 , for example by reaction with a suitable amine.

(Scheme 4)



All reactions for the synthesis of the compounds of the formula I are per se well-known to the skilled person and can be carried out under standard conditions according to or analogously to procedures described in the literature, for example in Houben-Weyl, Methoden der Organischen Chemie (Methods of Organic Chemistry), Thieme-Verlag, Stuttgart, or Organic Reactions, John Wiley & Sons, New York. Depending on the circumstances of the individual case, in order to avoid side reactions during the synthesis of a compound of the formula I it can be necessary or advantageous to temporarily block functional groups by introducing protective groups and to deprotect them in a later stage of the synthesis, or introduce functional groups in the form of precursor groups which in a later reaction step are converted into the desired functional groups. Such synthetic strategies and protective groups and precursor groups which are suitable in an individual case are known to the skilled person. If desired the compounds of the formula I can be purified by customary purification procedures, for example by recrystallization or chromatography. The starting compounds for the preparation of the compounds of the formula I are commercially available or can be prepared according to or analogously to literature procedures.

The compounds of the formula I according to the invention effect an increase of the cGMP concentration via the activation of the soluble guanylate cyclase (sGC), and they are therefore useful agents for the therapy and prophylaxis of disorders which are associated with a low or decreased cGMP level or which are caused thereby, or for

whose therapy or prophylaxis an increase of the present cGMP level is desired. The activation of the sGC by the compounds of the formula I can be examined, for example, in the activity assay described below.

5 Disorders and pathological conditions which are associated with a low cGMP level or in which an increase of the cGMP level is desired and for whose therapy and prophylaxis it is possible to use compounds of the formula I are, for example, cardiovascular diseases, such as endothelial dysfunction, diastolic dysfunction, atherosclerosis, hypertension, stable and unstable angina pectoris, thromboses, restenoses, myocardial
10 infarction, strokes, cardiac insufficiency or pulmonary hypertonia, or, for example, erectile dysfunction, asthma bronchiale, chronic kidney insufficiency and diabetes. Compounds of the formula I can additionally be used in the therapy of cirrhosis of the liver and also for improving a restricted memory performance or ability to learn.

15 The compounds of the formula I and their physiologically acceptable salts can be administered to animals, preferably to mammals, and in particular to humans, as pharmaceuticals by themselves, in mixtures with one another or in the form of pharmaceutical preparations. A subject of the present invention therefore also are the compounds of the formula I and their physiologically acceptable salts for use as
20 pharmaceuticals, their use for activating soluble guanylate cyclase, for normalizing a disturbed cGMP balance and in particular their use in the therapy and prophylaxis of the

abovementioned syndromes as well as their use for preparing medicaments for these purposes.

Furthermore, a subject of the present invention are pharmaceutical preparations (or
5 pharmaceutical compositions) which comprise as active component an effective dose of
at least one compound of the formula I and/or a physiologically acceptable salt thereof
and a customary pharmaceutically acceptable carrier, i. e. one or more
pharmaceutically acceptable carrier substances and/or additives. A subject of the
present invention is also those compounds of the formula I which were already known
per se and which are excluded by disclaimer from the above-defined compounds of the
40 formula I which are per se a subject of the present invention, and their physiologically
acceptable salt as activators of soluble guanylate cyclase.

All statements above and below relating to the pharmacological effects and the uses of
15 the compounds of the formula I thus also apply to the compound of the formula I
wherein simultaneously the ring A^2 which comprises the carbon atoms which carry the
groups $C(=X)-NH-$ and $NH-SO_2R^2$ is a benzene ring which is substituted in positions 3
and 5 by chlorine, R^2 is methyl, X is oxygen and $R^1-S(O)_n-A^1-$ is a 5-chloro-2-(4-
chlorophenylmercapto)-phenyl residue, and its physiologically acceptable salts.

Thus, a subject of the invention are, for example, said compound and its physiologically acceptable salts for use as a pharmaceutical, pharmaceutical preparations which comprise as active component an effective dose of said compound and/or a physiologically acceptable salt thereof and a customary pharmaceutically acceptable carrier, and the uses of said compound and/or a physiologically acceptable salt thereof in the therapy or prophylaxis of the abovementioned syndromes as well as their use for preparing medicaments for these purposes.

The pharmaceuticals according to the invention can be administered orally, for example in the form of pills, tablets, lacquered tablets, sugar-coated tablets, granules, hard and soft gelatin capsules, aqueous, alcoholic or oily solutions, syrups, emulsions or suspensions, or rectally, for example in the form of suppositories. Administration can also be carried out parenterally, for example subcutaneously, intramuscularly or intravenously in the form of solutions for injection or infusion. Other suitable administration forms are, for example, percutaneous or topical administration, for example in the form of ointments, tinctures, sprays or transdermal therapeutic systems, or the inhalative administration in the form of nasal sprays or aerosol mixtures, or, for example, microcapsules, implants or rods. The preferred administration form depends, for example, on the disease to be treated and on its severity.

The amount of active compound of the formula I and/or its physiologically acceptable salts in the pharmaceutical preparations normally is from 0.2 to 500 mg, preferably from 1 to 200 mg, per dose, but depending on the type of the pharmaceutical preparation it can also be higher. The pharmaceutical preparations usually comprise 0.5 to 90 percent
5 by weight of the compounds of the formula I and/or their physiologically acceptable salts. The preparation of the pharmaceutical preparations can be carried out in a manner known per se. For this purpose, one or more compounds of the formula I and/or their physiologically acceptable salts, together with one or more solid or liquid pharmaceutical carrier substances and/or additives (or auxiliary substances) and, if
10 desired, in combination with other pharmaceutically active compounds having therapeutic or prophylactic action, are brought into a suitable administration form or dosage form which can then be used as a pharmaceutical in human or veterinary medicine.

15 For the production of pills, tablets, sugar-coated tablets and hard gelatin capsules it is possible to use, for example, lactose, starch, for example maize starch, or starch derivatives, talc, stearic acid or its salts, etc. Carriers for soft gelatin capsules and suppositories are, for example, fats, waxes, semisolid and liquid polyols, natural or hardened oils, etc. Suitable carriers for the preparation of solutions, for example of
20 solutions for injection, or of emulsions or syrups are, for example, water, physiologically sodium chloride solution, alcohols such as ethanol, glycerol, polyols, sucrose, invert

sugar, glucose, mannitol, vegetable oils, etc. It is also possible to lyophilize the compounds of the formula I and their physiologically acceptable salts and to use the resulting lyophilisates, for example, for preparing preparations for injection or infusion. Suitable carriers for microcapsules, implants or rods are, for example, copolymers of

5 glycolic acid and lactic acid.

Besides the active compounds and carriers, the pharmaceutical preparations can also contain customary additives, for example fillers, disintegrants, binders, lubricants, wetting agents, stabilizers, emulsifiers, dispersants, preservatives, sweeteners,

10 colorants, flavorings, aromatizers, thickeners, diluents, buffer substances, solvents, solubilizers, agents for achieving a depot effect, salts for altering the osmotic pressure, coating agents or antioxidants.

The dosage of the active compound of the formula I to be administered and/or of a

15 physiologically acceptable salt thereof depends on the individual case and is, as is customary, to be adapted to the individual circumstances to achieve an optimum effect. Thus, it depends on the nature and the severity of the disorder to be treated, and also on the sex, age, weight and individual responsiveness of the human or animal to be treated, on the efficacy and duration of action of the compounds used, on whether the

20 therapy is acute or chronic or prophylactic, or on whether other active compounds are administered in addition to compounds of the formula I. In general, a daily dose of

approximately 0.01 to 100 mg/kg, preferably 0.1 to 10 mg/kg, in particular 0.3 to 5 mg/kg (in each case mg per kg of bodyweight) is appropriate for administration to an adult weighing approximately 75 kg in order to obtain the desired results. The daily dose can be administered in a single dose or, in particular when larger amounts are administered, be divided into several, for example two, three or four individual doses. In some cases, depending on the individual response, it may be necessary to deviate upwards or downwards from the given daily dose.

The compounds of the formula I activate the soluble guanylate cyclase, mainly by binding in the heme binding pocket of the enzyme. On account of this property, apart from use as pharmaceutically active compounds in human medicine and veterinary medicine, they can also be employed as a scientific tool or as aids for biochemical investigations in which such an effect on guanylate cyclase is intended, and also for diagnostic purposes, for example in the in vitro diagnosis of cell samples or tissue samples. The compounds of the formula I and salts thereof can furthermore be employed, as already mentioned above, as intermediates for the preparation of other pharmaceutically active compounds.

The following examples of compounds of the formula I and of intermediates for their preparation illustrate the invention without limiting it.

Examples

1.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoic acid

5 33.71 g (0.32 mol) of sodium carbonate were dissolved in 250 ml of water and warmed to 60 °C. 25.00 g (0.13 mol) of 2-amino-4,5-dimethoxy-benzoic acid were introduced into the solution, and to this solution 29.55 g (0.14 mol) of 4-chloro-benzenesulfonyl chloride were added portionwise over 15 min . After cooling the mixture was filtered with suction, the residue was taken up in 1 % sodium hydrogencarbonate solution, the
10 solution filtered, and the product precipitated by addition of 1 N hydrochloric acid. 25.90 g (55 %) of 2-(4-chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoic acid of melting point (m.p.) 212-214 °C were obtained.

Analogously were obtained:

15

2.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-benzoic acid; M.p.: 210 °C

3.) 5-Chloro-2-(3,4-dichloro-phenylsulfonylamino)-benzoic acid

20 5.) 2-(4-Chloro-phenylsulfonylamino)-cyclopentanecarboxylic acid; M.p.: 147 °C

6.) 2-(4-Chloro-phenylsulfonylamino)-5-methyl-benzoic acid; M.p.: 201 °C

7.) 3-(4-Chloro-phenylsulfonylamino)-thiophene-2-carboxylic acid; M.p.: 180 °C

5 8.) 3-(4-Chloro-phenylsulfonylamino)-pyrazole-4-carboxylic acid; Oil

9.) 2-(4-Chloro-phenylsulfonylamino)-pyridine-3-carboxylic acid; M.p.: Decomposition
(dec.) > 360 °C

10.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoyl chloride

25.90 g (0.07 mol) of 2-(4-chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoic acid were mixed with 75 ml of toluene, 17.30 g (0.08 mol) of phosphorus pentachloride were added and the mixture was stirred at 40-45 °C for 2.5 h. Then the mixture was concentrated in vacuo to half of its volume and the product that precipitated was filtered off with suction and washed with a small amount of toluene. 25.30 g (93 %) of 2-(4-chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoyl chloride having a melting point of 175-177 °C were obtained.

20 Analogously were obtained:

11.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-benzoyl chloride; M.p.: 127 °C

12.) 5-Chloro-2-(3,4-dichloro-phenylsulfonylamino)-benzoyl chloride; M.p.: 117 °C

5 13.) 2-(4-Chloro-phenylsulfonylamino)-cyclopentanecarboxylic acid chloride; M.p.: 107 °C

14.) 2-(4-Chloro-phenylsulfonylamino)-5-methyl-benzoyl chloride; M.p.: 114 °C

10 15.) 3-(4-Chloro-phenylsulfonylamino)-thiophene-2-carboxylic acid chloride; M.p.: 122 °C

16.) 3-(4-Chloro-phenylsulfonylamino)-pyrazole-4-carboxylic acid chloride; M.p.: 260 °C
(Dec.)

15

17.) 2-(4-Chloro-phenylsulfonylamino)-pyridine-3-carboxylic acid chloride

18.) 4-((2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoyl)-amino)-
benzenesulfonyl fluoride

20

10.00 g (25.6 mmol) of 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoyl chloride were mixed with 300 ml of toluene, 4.49 g (25.6 mmol) of 4-aminobenzenesulfonyl fluoride were added and the mixture was heated under reflux for 4 h. After cooling the precipitated solid was filtered off with suction and washed with
5 toluene. 11.71 g (87 %) of the title compound having a melting point of 216-219 °C were obtained.

Analogously were obtained:

19.) 4-((5-Chloro-2-(4-chloro-phenylsulfonylamino)-benzoyl)-amino)-benzenesulfonyl fluoride; M.p.: 242 °C

20.) N-(4-Aminosulfonyl-phenyl)-5-chloro-2-(4-chloro-phenylsulfonylamino)-benzamide; M.p.: 260 °C

21.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-((4-(4-nitro-phenyl)-mercapto)-phenyl)-benzamide; M.p.: 255 °C

22.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(phenylmercapto)-phenyl)-benzamide; M.p.: 169 °C

23.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-methylmercapto-phenyl)-
benzamide; M.p.: 220 °C

24.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(2-methyl-benzothiazol-5-yl)-

5 benzamide; M.p.: 251 °C

25.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(3-diethylamino-2-hydroxy-propyl-
mercapto)-phenyl)-benzamide; M.p.: 102 °C

26.) 4-((5-Chloro-2-(3,4-dichloro-phenylsulfonylamino)-benzoyl)-amino)-
benzenesulfonyl fluoride; M.p.: 232 °C

27.) 4-(2-(4-Chloro-phenylsulfonylamino)-cyclopentanecarbonylamino)-benzenesulfonyl
fluoride; M.p.: 211 °C

28.) 4-((2-(4-Chloro-phenylsulfonylamino)-5-methyl-benzoyl)-amino)-benzenesulfonyl
fluoride; M.p.: 224 °C

29.) 4-((3-(4-Chloro-phenylsulfonylamino)-thiophene-2-carbonyl)-amino)-

20 benzenesulfonyl fluoride; M.p.: 255 °C

30.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-mercapto-phenyl)-benzamide;

M.p.: 202 °C

31.) 4-((3-(4-Chloro-phenylsulfonylamino)-pyrazol-4-carbonyl)-amino)-benzenesulfonyl

5 fluoride; M.p.: 251 °C

32.) 3-((5-Chloro-2-(4-chloro-phenylsulfonylamino)-benzoyl)-amino)-benzenesulfonyl

fluoride; M.p.: 224 °C

33.) 4-(2-(4-Chloro-phenylsulfonylamino)-pyridine-3-carbonyl)-amino)-benzenesulfonyl

fluoride; M.p.: 263-265 °C

34.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-methyl-5-(thiomorpholine-4-

sulfonyl)-thiazol-2-yl)-benzamide; M.p.: 265-267 °C

35.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(2-methylmercapto-phenyl)-

benzamide

36.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(3-methylmercapto-phenyl)-

benzamide

37.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(5-methyl-isoxazol-3-yl-sulfamoyl)-phenyl)-benzamide

38.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(4-nitro-phenylsulfonyl)-phenyl)-

5 benzamide

39.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(5-ethylsulfonyl-2-hydroxy-phenyl)-benzamide

40.) N-(3-Butylsulfamoyl-phenyl)-5-chloro-2-(4-chloro-phenylsulfonylamino)-benzamide

41.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(2-nitro-5-propylmercapto-phenyl)-benzamide

42.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-thiocyanato-phenyl)-benzamide

43.) N-(4-Acetylsulfamoyl-phenyl)-5-chloro-2-(4-chloro-phenylsulfonylamino)-benzamide

44.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(2-phenylmercapto-phenyl)-benzamide

45.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(2-chloro-5-(2-cyano-ethylsulfamoyl)-phenyl)-benzamide

46.) N-(5-Butylsulfamoyl-2-methoxy-phenyl)-5-chloro-2-(4-chloro-phenylsulfonylamino)-

5 benzamide

47.) N-(4-Benzoylsulfamoyl-phenyl)-5-chloro-2-(4-chloro-phenylsulfonylamino)-benzamide

48.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(2-chloro-4-methylsulfonyl-phenyl)-benzamide

49.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(hexadecylsulfonyl)-phenyl)-benzamide

15

50.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(butylaminocarbonylamino-sulfonyl)-phenyl)-benzamide

51.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(2-sulfamoyl-phenyl)-benzamide

20

52.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(2-methylmercapto-5-trifluoromethyl-phenyl)-benzamide

53.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(3-methylsulfonyl-phenyl)-benzamide

5

54.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(3-(2-hydroxy-ethylsulfonyl)-phenyl)-benzamide

55.) (4-(5-Chloro-2-(4-chloro-phenylsulfonylamino)-benzoylamino)-phenylmercapto)-acetic acid

10

56.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(3,4-dimethyl-isoxazol-5-ylsulfamoyl)-phenyl)-benzamide

15 57.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(thiazol-2-ylsulfamoyl)-phenyl)-benzamide

58.) 5-Chloro-2-(3,4-dichloro-phenylsulfonylamino)-N-(4-ethylmercapto-phenyl)-benzamide; M.p.: 171 °C

20

59.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide

500 mg (0.95 mmol) of 4-((2-(4-chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoyl)-amino)-benzenesulfonyl fluoride were dissolved in 1 ml of thiomorpholine and heated to 90 °C for 30 min. For working up the mixture was poured onto 50 ml of ice/1 N hydrochloric acid, the precipitate was filtered off with suction, dried in a vacuum-drying chamber over phosphorus pentoxide and recrystallized from hexane/ethyl acetate. 378 mg (65 %) of the title compound having a melting point of 241 °C were obtained.

Analogously were obtained:

60.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(cis-2,6-dimethyl-morpholine-4-sulfonyl)-phenyl)-pyridine-3-carboxamide; M.p.: 256-258 °C

61.) N-(4-(4-Carbamoyl-piperidine-1-sulfonyl)-phenyl)-2-(4-chloro-phenylsulfonylamino)-pyridine-3-carboxamide; M.p.: 273-276 °C

62.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(piperidine-1-sulfonyl)-phenyl)-pyridine-3-carboxamide; M.p.: 180-183 °C

63.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 246 °C

64.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(4-methyl-piperazine-1-sulfonyl)-phenyl)-benzamide; M.p.: 219 °C

65.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(2,6-dimethyl-morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 259 °C

66.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(cis-2,6-dimethyl-morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 251 °C

67.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(4-hydroxy-piperidine-1-sulfonyl)-phenyl)-benzamide; M.p.: 255 °C

68.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(1,4-dioxo-8-aza-spiro[4.5]decane-8-sulfonyl)-phenyl)-benzamide; M.p.: 256 °C

69.) 5-Chloro-2-(3,4-dichloro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 253 °C

70.) 5-Chloro-2-(3,4-dichloro-phenylsulfonylamino)-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 222 °C

71.) 5-Chloro-2-(3,4-dichloro-phenylsulfonylamino)-N-(4-(4-methyl-piperazine-1-sulfonyl)-phenyl)-benzamide; M.p.: 246 °C

72.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 172 °C

73.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(4-(2-hydroxy-ethyl)-piperazine-1-sulfonyl)-phenyl)-benzamide; M.p.: 277 °C

74.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-cyclopentanecarboxamide; M.p.: 180 °C

75.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-diethylsulfamoyl-phenyl)-benzamide; M.p.: 226 °C

76.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(piperidine-1-sulfonyl)-phenyl)-benzamide; M.p.: 240 °C

77.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(2-methoxy-ethylsulfamoyl)-phenyl)-benzamide; M.p.: 209 °C

78.) 2-(4-Chloro-phenylsulfonylamino)-5-methyl-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 203 °C

79.) 3-(4-Chloro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-thiophene-2-carboxamide; M.p.: 220 °C

80.) 3-(4-Chloro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-1H-pyrazole-4-carboxamide; Oil

81.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(3-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 238 °C

82.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(3-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 202 °C

83.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(3-(4-methyl-piperazine-1-sulfonyl)-phenyl)-benzamide hydrochloride; M.p.: 245 °C

84.) 3-(4-Chloro-phenylsulfonylamino)-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-
thiophene-2-carboxamide; M.p.: 229 °C

85.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-

5 benzamide; M.p.: 228 °C

86.) 2-(4-Chloro-phenylsulfonylamino)-5-methyl-N-(4-(thiomorpholine-4-sulfonyl)-
phenyl)-benzamide; M.p.: 234 °C

87.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(4-methyl-piperazine-1-
sulfonyl)-phenyl)-benzamide; M.p.: 172 °C

88.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(cis-2,6-dimethyl-morpholine-4-sulfonyl)-
phenyl)-4,5-dimethoxy-benzamide; M.p.: 208 °C

89.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(4-hydroxy-piperidine-1-sulfonyl)-phenyl)-
4,5-dimethoxy-benzamide; M.p.: 244 °C

90.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(piperidine-3-sulfonyl)-

phenyl)-benzamide; M.p.: 258 °C

91.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(thiazolidine-3-sulfonyl)-phenyl)-benzamide; M.p.: 261 °C

92.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(2,5-dihydro-1H-pyrrole-1-sulfonyl)-phenyl)-4,5-dimethoxy-benzamide; M.p.: 262 °C

93.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(1,2,3,6-tetrahydro-pyridine-1-sulfonyl)-phenyl)-4,5-dimethoxy-benzamide; M.p.: 252 °C

94.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(2-methyl-piperidine-1-sulfonyl)-phenyl)-benzamide; M.p.: 227 °C

95.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(piperazine-1-sulfonyl)-phenyl)-benzamide; M.p.: 243 °C

96.) 4-(4-(2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoylamino)-phenylsulfonyl)-piperazine-1-carboxylic acid ethyl ester; M.p.: 245 °C

97.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(4-methyl-piperidine-1-sulfonyl)-phenyl)-benzamide; M.p.: 267 °C

98.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(4-methyl-perhydro-[1,4]diazepine-1-sulfonyl)-phenyl)-benzamide; M.p.: 274 °C

99.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(4-ethyl-piperazine-1-sulfonyl)-phenyl)-4,5-dimethoxy-benzamide; M.p.: 191 °C

100.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-((2-dimethylamino-ethyl)-ethyl-sulfamoyl)-phenyl)-4,5-dimethoxy-benzamide; M.p.: Dec. > 119 °C

101.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(1,4,5,6-tetrahydro-pyrimidine-1-sulfonyl)-phenyl)-4,5-dimethoxy-benzamide; M.p.: Dec. > 237 °C

102.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(4-(pyrimidin-2-yl)-piperazine-1-sulfonyl)-phenyl)-benzamide; M.p.: Dec. > 194 °C

103.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(4-(4-chloro-phenyl)-piperazine-1-sulfonyl)-phenyl)-4,5-dimethoxy-benzamide; M.p.: Dec. > 243 °C

104.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(indan-1-ylsulfamoyl)-phenyl)-4,5-dimethoxy-benzamide; M.p.: 161 °C

105.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-((2-(1H-indol-3-yl)-ethyl)-methyl-sulfamoyl)-phenyl)-4,5-dimethoxy-benzamide; M.p.: 182 °C

106.) 1-(4-((2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoyl)-amino)-phenylsulfonyl)-piperidine-4-carboxamide; M.p.: 252 °C

107.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-cyclopropylsulfamoyl-phenyl)-4,5-dimethoxy-benzamide; M.p.: 222 °C

108.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(3-hydroxy-pyrrolidine-1-sulfonyl)-phenyl)-4,5-dimethoxy-benzamide; M.p.: 272 °C

109.) N-(4-(Allyl-cyclohexyl-sulfamoyl)-phenyl)-2-(4-chloro-phenylsulfonylamino)-4,5-dimethoxy-benzamide; M.p.: 182 °C

110.) 1-(4-((2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-benzoyl)-amino)-phenylsulfonyl)-pyrrolidine-2-carboxylic acid; M.p.: 240 °C (sintering)

111.) 5-Chloro-2-nitro-benzoyl chloride

100.00 g (0.50 mol) of 5-chloro-2-nitrobenzoic acid were mixed with 72.20 g (0.61 mol) of thionyl chloride and the mixture was heated under reflux for 2 h. The excess thionyl chloride was removed in vacuo. 106.50 g (ca. 98 %) of crude 5-chloro-2-nitro-benzoyl chloride were obtained as an oil.

5

Analogously was obtained:

112.) 5-Methyl-2-nitro-benzoyl chloride; Oil

113.) 4-(5-Chloro-2-nitro-benzoylamino)-benzenesulfonyl fluoride

86.00 g (0.39 mol) of 5-chloro-2-nitro-benzoyl chloride was dissolved in 300 ml of toluene, a solution of 62.00 g (0.35 mol) of 4-aminobenzenesulfonyl fluoride was added dropwise, and the mixture was heated under reflux for 4 h. Subsequently it was cooled, concentrated in vacuo to half of its volume, cooled, and the precipitated solid was filtered off with suction. 121.60 g (86 %) of the title compound having a melting point of 182-184 °C were obtained.

Analogously were obtained:

114.) 4-(5-Methyl-2-nitro-benzoylamino)-benzenesulfonyl fluoride; M.p.: 179 °C

115.) 5-Chloro-N-(4-ethylmercapto-phenyl)-2-nitro-benzamide

116.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-nitro-benzamide

5 120.00 g (0.33 mol) of 4-(5-chloro-2-nitro-benzoylamino)-benzenesulfonyl fluoride, 29.10 g (0.33 mol) of morpholine and 33.85 g (0.33 mol) of triethylamine were stirred in 1200 ml of toluene at 60 °C for 2 days. The precipitated solid was filtered off with suction and recrystallized from acetone/n-hexane. 102.10 g (71 %) of the title compound having a melting point of 243-245 °C were obtained.

Analogously were obtained:

117.) 5-Chloro-2-nitro-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 120 °C

118.) 5-Methyl-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-nitro-benzamide; M.p.: 249 °C

119.) N-(4-(Morpholine-4-sulfonyl)-phenyl)-5-(morpholin-4-yl)-2-nitro-benzamide

20 20.00 g (0.56 mol) of 4-(5-chloro-2-nitro-benzoylamino)-benzenesulfonyl fluoride in 48.5 g (0.557 mol) of morpholine were heated under reflux for 1 h. Subsequently the mixture

was cooled, poured onto ice/hydrochloric acid and filtered with suction. 26.0 g (98 %) of the title compound having a melting point of 252 °C were obtained.

120.) 2-Amino-5-chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide

5

11.10 g (26.1 mmol) of 5-chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-nitro-benzamide were dissolved in 440 ml of tetrahydrofuran/methanol (1:1) and a solution of 27.23 g (156.4 mmol) of sodium dithionite in 330 ml of water was added dropwise. After stirring for 1 h at room temperature the organic solvents were removed in rotary evaporator, and the precipitated product was filtered off with suction and purified by chromatography over silica with methylene chloride/methanol (9:1). 5.68 g (55 %) of the title compound having a melting point of 229-231 °C were obtained.

Analogously were obtained:

15

121.) 2-Amino-5-chloro-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 177 °C

122.) 2-Amino-N-(4-(morpholine-4-sulfonyl)-phenyl)-(5-morpholin-4-yl)-benzamide;

20 M.p.: 228 °C

123.) 2-Amino-5-chloro-N-(4-ethylsulfonyl-phenyl)-benzamide; M.p.: 159-161 °C

124.) 5-Chloro-2-(5-chloro-1,3-dimethyl-1H-pyrazole-4-sulfonyl-amino)-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide

5

250 mg (0.60 mmol) of 2-amino-5-chloro-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide were dissolved in 10 ml of dry pyridine, and a solution of 195 mg (0.85 mmol) 5-chloro-1,3-dimethyl-1H-pyrazole-4-sulfonyl chloride in 5 ml of pyridine was added dropwise at 0 °C. After 2 h the mixture was poured onto ice, the precipitated solid was filtered off with suction and purified by chromatography over silica with methylene chloride/methanol (98:2). 250 mg (69 %) of the title compound having a melting point of 215-216 °C were obtained.

Analogously were obtained:

15

125.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(4-methyl-phenylsulfonylamino)-benzamide; M.p.: 214 °C

20 126.) 5-Chloro-2-(3,4-dimethoxy-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 245 °C

127.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(4-trifluoromethoxy-phenylsulfonylamino)-benzamide; M.p.: 195 °C

128.) 2-((4-Acetylamino-phenyl)-sulfonylamino)-5-chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 198 °C

129.) 5-Chloro-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 112 °C

130.) 5-Chloro-2-(5-chloro-1,3-dimethyl-pyrazole-4-sulfonyl-amino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 161 °C

131.) 5-Chloro-2-((1-methyl-imidazole-4-sulfonyl)-amino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 141 °C

132.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(pyridine-3-sulfonylamino)-benzamide; M.p.: 222 °C

133.) 2-(4-Benzoyloxy-phenylsulfonylamino)-5-chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 245 °C

134.) 5-Chloro-2-(ethylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide;
M.p.: 274-276 °C

135.) 2-((2-Acetamideo-4-methyl-thiazole-5-sulfonyl)-amino)-5-chloro-N-(4-(morpholine-
5 4-sulfonyl)-phenyl)-benzamide; M.p.: 257 °C

136.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(thiophene-2-sulfonylamino)-
benzamide; M.p.: 216 °C

137.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(4-trifluoromethyl-
10 phenylsulfonylamino)-benzamide; M.p.: 264 °C

138.) 2-(4-Bromo-phenylsulfonylamino)-5-chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-
benzamide; M.p.: 232 °C

15

139.) 2-(3,5-Bis-trifluoromethyl-phenylsulfonylamino)-5-chloro-N-(4-(morpholine-4-
sulfonyl)-phenyl)-benzamide; M.p.: 209 °C

140.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(4-nitro-phenylsulfonylamino)-
20 benzamide; M.p.: 239 °C

141.) 5-Chloro-2-(4-cyano-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 238 °C

142.) 5-Chloro-2-(4-methylsulfonyl-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 181 °C

143.) 5-Chloro-2-(4-isopropyl-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 105 °C

144.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-((2-phenyl-ethenyl)-sulfonylamino)-benzamide; M.p.: 278 °C

145.) 5-Chloro-2-(4,5-dibromo-thiophene-2-sulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 232 °C

146.) 5-Chloro-2-(4-fluoro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 245 °C

147.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(5-phenylsulfonyl-thiophene-2-sulfonylamino)-benzamide; M.p.: 103 °C

148.) 5-Chloro-2-(3-chloro-4-methoxy-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 274 °C

149.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(quinoline-8-sulfonylamino)-

5 benzamide; M.p.: 262 °C

150.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(2,4,6-trimethyl-phenylsulfonylamino)-benzamide; M.p.: 240 °C

10 151.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(3-nitro-phenylsulfonylamino)-benzamide; M.p.: 220 °C

152.) 5-Chloro-2-(4-methoxy-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 269 °C

15

153.) 5-Chloro-2-methylsulfonylamino-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 248 °C

154.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-phenylmethylsulfonylamino-

20 benzamide; M.p.: 106 °C

155.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(2,2,2-trifluoro-ethylsulfonylamino)-benzamide; M.p.: 208 °C

156.) 2-(Butyl-sulfonylamino)-5-chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide;

5 M.p.: 102 °C

157.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(3-trifluoromethyl-phenylsulfonylamino)-benzamide; M.p.: 212 °C

158.) 2-(4-Bromo-2,5-dichloro-thiophene-3-sulfonylamino)-5-chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 267 °C

159.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(2-trifluoromethyl-phenylsulfonylamino)-benzamide; M.p.: 234 °C

160.) 5-Chloro-2-(3-chloro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 206 °C

161.) 2-(4-Bromo-2-methoxy-phenylsulfonylamino)-5-chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 260 °C

162.) 5-Chloro-2-(2,6-dichloro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 244 °C

163.) 5-Chloro-2-(2-cyano-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 200 °C

164.) 2-(4-Butoxy-phenylsulfonylamino)-5-chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 225 °C

165.) 5-Chloro-2-(7,7-dimethyl-2-oxo-bicyclo[2.2.1]heptane-1-sulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 120 °C

166.) 5-Chloro-2-(3-fluoro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 204 °C

167.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-5-(morpholin-4-yl)-benzamide; M.p.: 264 °C

168.) 5-Chloro-N-(4-ethylsulfonyl-phenyl)-2-(4-methyl-phenylsulfonylamino)-benzamide; M.p.: 188-192 °C

169.) 5-Chloro-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-ethylsulfonyl-phenyl)-benzamide; M.p.: 195-197 °C

170.) 5-Chloro-2-(4-chloro-3-nitro-phenylsulfonylamino)-N-(4-ethylsulfonyl-phenyl)-

5 benzamide; M.p.: 196-198 °C

171.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-ethylsulfonyl-phenyl)-benzamide; M.p.: 180-185 °C

172.) 5-Chloro-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(4-ethylsulfonyl-phenyl)-benzamide; M.p.: Dec. > 249 °C

173.) 5-Chloro-2-ethylsulfonylamino-N-(4-ethylsulfonyl-phenyl)-benzamide; M.p.: 103 °C

174.) 4-Chloro-N-(2-(1H-benzimidazol-2-yl)-4-chloro-phenyl)-benzenesulfonamide

1.00 g (2.7 mmol) of 5-chloro-2-(4-chloro-phenylsulfonylamino)-benzoyl chloride and 296 mg (2.7 mmol) of o-phenylenediamine in 150 ml of toluene were heated under reflux for 1 h. A small amount of solid was filtered off with suction and the filtrate was evaporated. The residue was taken up in 50 ml of toluene, 600 mg of thionyl chloride

were added and the mixture was again heated under reflux for 10 h. Subsequently it was cooled and the precipitated solid was filtered off with suction. 280 mg (25 %) of the title compound having a melting point of 225-228 °C were obtained.

- 5 175.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(1,1-dioxo-thiomorpholine-4-sulfonyl)-phenyl)-benzamide

and

- 176.) 2-(4-Chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(1-oxo-thiomorpholine-4-sulfonyl)-phenyl)-benzamide

500 mg (0.82 mmol) of 2-(4-chloro-phenylsulfonylamino)-4,5-dimethoxy-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide in 50 ml of acetone were cooled to 0 °C. A solution of 371 mg (1.23 mmol) of 57 % m-chloroperbenzoic acid in 20 ml of acetone was added and the mixture was stirred at room temperature over night. For working up
15 it was poured onto water/ice and the precipitate was filtered off with suction. The two products obtained as a mixture were separated by chromatography over silica with methylene chloride/methanol (97:3).

Analogously were obtained:

177.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(1,1-dioxo-thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 182 °C

178.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(1-oxo-thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 233 °C

179.) 5-Chloro-2-(3,4-dichloro-phenylsulfonylamino)-N-(4-ethylsulfonyl-phenyl)-benzamide; M.p.: 240 °C

180.) 5-Chloro-N-(4-ethylsulfonyl-phenyl)-2-nitro-benzamide

181.) 4-Chloro-N-(4-chloro-2-(6-(morpholine-4-sulfonyl)-1H-benzimidazol-2-yl)-phenyl)-benzenesulfonamide

200 mg (0.5 mmol) of 4-chloro-N-(2-(1H-benzimidazol-2-yl)-4-chloro-phenyl)-benzenesulfonamide were added at 0 °C to 1 ml of chlorosulfuric acid and heated to 60 °C for 30 min. Subsequently the mixture was poured onto water/ice, and the solid was filtered off with suction, dried and added at 0 °C to 1 ml of morpholine. After stirring at room temperature for 1 h the mixture was poured onto ice/hydrochloric acid and extracted with ethyl acetate. The extracts were evaporated and the residue was purified

by chromatography over silica with hexane/ethyl acetate (1:1). 20 mg (7 %) of the title compound having a melting point of 225-228 °C were obtained.

¹H-NMR (D₆-DMSO): δ (ppm) = 2.9 (m, 4 H, morpholine-H), 3.6 (m, 4 H, morpholine-H), 7.5 (dd, 4 H, phenylene-H), 7.4-8.2 (m, 6 H, benzo-H, phenyl-H)

5

182.) 5-Chloro-N-(4-(morpholine-4-sulfonyl)-phenyl)-2-(2-(pyrrolidin-1-yl)-ethylsulfonylamino)-benzamide

The compound was prepared by using 2-chloro-ethylsulfonyl chloride. The 1-(2-(4-chloro-2-(4-(morpholine-4-sulfonyl)-phenylcarbamoyl)-phenylsulfamoyl)-ethyl)-pyridinium chloride that was isolated as an intermediate was reacted with pyrrolidine to give the title compound.

¹H-NMR (D₆-DMSO): δ (ppm) = 1.8 (m, 4 H, pyrrolidine-H), 2.65 (m, 4 H, pyrrolidine-H), 3.0 (m, 4 H, morpholine-H), 3.1 (t, 2 H, ethylene-H), 3.35 (t, 2 H, ethylene-H), 3.75 (m, 4 H, morpholine-H), 7.50 (dd, 1 H, H-4), 7.7 (d, 1 H, H-3), 7.75 (dd, 1 H, H-6), 7.85 ("dd", 4 H, phenylene-H)

183.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-isopropylmercapto-phenyl)-benzamide

1.00 g (2.21 mmol) of 5-chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-mercapto-phenyl)-benzamide were dissolved in 25 ml of dimethylformamide and 0.25 g (2.21 mmol) of potassium tert-butyrate were added. The mixture was stirred at room temperature for 15 min, then 0.27 g (2.21 mmol) of isopropyl bromide were added dropwise, and the mixture was heated to 60 °C for 8 h. For working up it was poured onto water and extracted with ethyl acetate. The combined extracts were evaporated and the residue was purified by chromatography over silica with hexane/ethyl acetate (3:1). 420 mg (39 %) of the title compound having a melting point of 168-169 °C were obtained.

Analogously were obtained:

184.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-cyanomethylmercapto-phenyl)-benzamide; M.p.: 104 °C

185.) (4-((5-Chloro-2-(4-chloro-phenylsulfonylamino)-benzoyl)-amino)-phenylmercapto)-acetic acid ethyl ester; M.p.: 133 °C

186.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(2-(morpholin-4-yl)-ethylmercapto)-phenyl)-benzamide; M.p.: 95 °C

187.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(2-(2-methoxy-ethoxy)-ethylmercapto)-phenyl)-benzamide; Oil

188.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(prop-2-ynyl)-mercapto-phenyl)-
5 benzamide; M.p.: 185 °C

189.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-isopropylmercapto-phenyl)-
benzamide; M.p.: 169 °C

10 190.) 5-Chloro-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(morpholine-4-sulfonyl)-
phenyl)-benzamide sodium salt

A mixture of 0.48 g finely powdered sodium hydroxide and 7 g of 5-chloro-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide in 250 ml
15 of ethanol was brought into solution by short heating. Then the mixture was evaporated in vacuo, 50 ml of water were added and it was again evaporated in vacuo to dryness. This procedure was repeated twice. The resulting product was dried in vacuo at 50 °C. M.p.: 343 °C (Dec.)

20 Analogously to the above compounds the following example compounds were obtained:

191.) 4,5-Dimethoxy-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(N-methyl-N-(pyridin-3-yl-methyl)-aminosulfonyl)-phenyl)-benzamide hydrochloride; M.p.: 214°C

192.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(morpholine-4-sulfonyl)-3-methyl-phenyl)-benzamide; M.p.: 192 °C

193.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(cis-2,6-dimethyl-morpholine-4-sulfonyl)-3-methyl-phenyl)-benzamide; M.p.: 254 °C

194.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(3,5-dimethyl-piperidine-1-sulfonyl)-3-methyl-phenyl)-benzamide; M.p.: 242 °C

195.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(piperidine-1-sulfonyl)-3-methyl-phenyl)-benzamide; M.p.: 189 °C

196.) 4,5-Dimethoxy-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(4-(N-methyl-N-(pyridin-3-yl-methyl)-aminosulfonyl)-phenyl)-benzamide; M.p.: 213 °C

197.) 4,5-Dimethoxy-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(4-(N-(pyridin-3-yl-methyl)-aminosulfonyl)-phenyl)-benzamide; M.p.: 216 °C

198.) 5-Chloro-2-(2,4-dimethyl-thiazole-5-sulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 190 °C

199.) 4,5-Dimethoxy-2-(4-chloro-phenylsulfonylamino)-N-(4-(3,5-dimethyl-piperidine-1-sulfonyl)-phenyl)-benzamide; M.p.: 249 °C (Dec.)

200.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-(N-methyl-N-(pyridin-3-yl-methyl)-aminosulfonyl)-phenyl)-benzamide; Resin

201.) 3,4-Dimethoxy-2-(4-chloro-phenylsulfonylamino)-N-(4-(N-methyl-N-(pyridin-3-yl-methyl)-aminosulfonyl)-phenyl)-benzamide; M.p.: 241 °C

202.) 5-Bromo-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(morpholine-4-sulfonyl)-3-methyl-phenyl)-benzamide; M.p.: 249 °C

203.) 5-Bromo-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(cis-2,6-dimethyl-morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 244 °C

204.) 5-Bromo-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 197 °C

205.) 4,5-Dimethoxy-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(1,2,3,4-tetrahydro-isoquinoline-2-sulfonyl)-phenyl)-benzamide; M.p.: 213 °C

206.) 4,5-Dimethoxy-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 232 °C

207.) 4,5-Dimethoxy-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(cis-2,6-dimethyl-piperidine-1-sulfonyl)-phenyl)-benzamide; M.p.: 213 °C

208.) 5-Chloro-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(4-(1,2,3,4-tetrahydro-isoquinoline-2-sulfonyl)-phenyl)-benzamide; M.p.: 260 °C

209.) 5-Chloro-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(4-(N-methyl-N-(pyridin-3-yl-methyl)-aminosulfonyl)-phenyl)-benzamide; M.p.: 65 °C (sintering)

210.) 6-Methyl-2-(4-chloro-phenylsulfonylamino)-N-(4-(perhydroazepine-1-sulfonyl)-phenyl)-benzamide; M.p.: 151 °C

211.) 6-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(pyrrolidine-1-sulfonyl)-phenyl)-benzamide; M.p.: 217 °C

212.) 6-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(4-hydroxybutylamino)-sulfonyl)-phenyl)-benzamide; Resin

213.) 5-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(N-ethyl-N-(pyridin-4-yl-methyl)-aminosulfonyl)-phenyl)-benzamide; Resin

214.) 2-(4-Chloro-phenylsulfonylamino)-N-(4-thiomorpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 209 °C

215.) 3-Methyl-2-(4-chloro-phenylsulfonylamino)-N-(4-(N-methyl-N-(2-(pyridin-2-yl)-ethyl)-aminosulfonyl)-phenyl)-benzamide; M.p.: 193 °C

216.) 4,5-Difluoro-2-(4-chloro-phenylsulfonylamino)-N-(4-(4-aminocarbonyl-piperidine-1-sulfonyl)-phenyl)-benzamide; M.p.: 227 °C

217.) 4,5-Difluoro-2-(4-chloro-phenylsulfonylamino)-N-(4-(4-(2-hydroxyethyl)-piperazine-1-sulfonyl)-phenyl)-benzamide; Resin

218.) 5-Chloro-4-methoxy-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; Oil

219.) 5-Chloro-4-methoxy-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(cis-2,6-dimethyl-morpholine-1-sulfonyl)-phenyl)-benzamide; M.p.: 89 °C

220.) 5-Chloro-4-methoxy-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-(N-pyridin-3-yl)-N-methyl-aminosulfonyl)-phenyl)-benzamide; M.p.: 135 °C

221.) 4,5-Dimethoxy-2-(4-chloro-phenylsulfonylamino)-N-(4-(cis-2,6-dimethyl-morpholine-4-sulfonyl)-phenyl)-benzamide sodium salt; M.p.: 330 °C (Dec)

222.) 5-Chloro-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(4-(cis-2,6-dimethyl-morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 230 °C

223.) 5-Chloro-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(4-(3,5-dimethylpiperidine-1-sulfonyl)-phenyl)-benzamide; M.p.: 61 °C

224.) 5-Chloro-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(4-(morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 286 °C

225.) 5-Chloro-2-(5-chloro-thiophene-2-sulfonylamino)-N-(4-phenylsulfonyl)-phenyl-benzamide; M.p.: 227 °C

226.) 4-Chloro-2-(4-chloro-phenylsulfonylamino)-N-(4-(cis-2,6-dimethyl-morpholine-4-sulfonyl)-phenyl)-benzamide; M.p.: 103 °C.

Pharmacological investigations

(1) Activation of the soluble guanylate cyclase

The activation of the soluble guanylate cyclase (sGC) which catalyzes the conversion of guanosine triphosphate (GTP) into cyclic guanosine monophosphate (cGMP) and pyrophosphate, by the compounds according to the invention was quantified with the aid of an enzyme immunoassay (EIA) from Amersham. For this purpose the substances to be tested were initially incubated with sGC in microtiter plates, and the amount of the cGMP formed was then determined.

The sGC which was employed had been isolated from bovine lung (see Methods in Enzymology, Volume 195, p. 377). The test solutions (100 µl per well) contained 50 mM triethanolamine (TEA) buffer (pH 7.5), 3 mM MgCl₂, 3 mM reduced glutathione (GSH), 0.1 mM GTP, 1 mM 3-isobutyl-1-methylxanthine (IBMX), suitably diluted enzyme solution and the substance to be tested or, in the control experiments, the solvent. The substances to be tested were dissolved in dimethyl sulfoxide (DMSO) and the solution was diluted with DMSO/water, so that the final concentration (c) of the substance to be tested in the test solution had the value indicated in the table. The DMSO concentration in the test solution was 5 % (v/v). The reaction was initiated by addition of the sGC. The

reaction mixture was incubated at 37 °C for 15 to 20 minutes and then stopped by ice-cooling and addition of the stop reagent (50 mM EDTA, pH 8.0). An aliquot of 50 µl was taken and used for determining the cGMP content using the acetylation protocol of the Amersham cGMP-EIA kit. The absorption of the samples was measured at 450 nm (reference wavelength 620 nm) in a microtiter plate reader. The cGMP concentration was determined using a standard curve which was obtained under the same test conditions. The activation of sGC by a test substance is given as the n-fold stimulation of the basal enzyme activity which was found in the control experiments (using solvent instead of test substance) (calculated using the formula

$$\text{n-fold stimulation} = [\text{cGMP}]_{\text{test substance}} / [\text{cGMP}]_{\text{control}}).$$

The following results were obtained:

Compound of	Concentration "c"	"n-fold" stimulation
Example No.	[µM]	
23	50	14.7
59	50	34.8
63	50	33.9
64	50	23.9
65	50	24.6
66	50	33
67	50	29.6
68	50	12.1
69	50	28.3
70	50	25.1
71	50	13.4
72	50	27
73	50	16.5
75	50	5.1
76	50	10.6
77	50	5.9
79	50	15.4
84	50	23.7

	Compound of	Concentration "c"	"n-fold" stimulation
	Example No.	[μ M]	
	86	50	32.9
	87	50	12.5
	88	50	24.4
	89	50	11.6
5	124	50	31.2
	129	50	8.6
	130	50	35.3
	132	50	9.9
	134	50	7.2
10	136	50	24.2
	137	50	4.6
	139	50	21.9
	145	50	8
	146	50	10.2
15	148	50	15.5
	150	50	15.3
	151	50	19.9
	155	50	7.8
	156	50	7.8
20	157	50	4.6
	175	50	21.1
	176	50	13.9
	191	10	27.5
	192	10	26.7
25	193	10	31.1
	194	5	20.0
	195	10	16.2
	196	10	21.9
	206	50	19.3
30	208	10	23.1
	209	10	28.5
	222	10	29.5
	223	10	27.1
	224	50	27.8
35	225	10	13.4
	226	25	3.3

(2) Relaxation of rat aorta

For this test, normotensive male Wistar-Kyoto rats were sacrificed by a blow to the neck.

5 The abdominal cavity and the thorax were opened by a medium sternotomy. The descending aorta was subsequently removed, freed of connective tissue and divided into 8 rings of a length of approximately 4 mm. The tip of a pair of tweezers was introduced into the lumen of 4 of the 8 rings. The endothelium was removed by carefully rolling the rings over the tip of the pair of tweezers. All 8 aorta rings (4 with endothelium and 4 without endothelium) were subsequently suspended in an organ bath (Schuler-Organbad; Hugo Sachs Elektronik) at a constant temperature of 37 °C for the isometric determination of the contractile tone. For 30 minutes, the rings were calibrated at a resting tension of 1 g in carbonated (95 % O₂; 5 % CO₂) Krebs-Henseleit solution (composition: Na⁺ 144.0 mM; K⁺ 5.9 mM; Cl⁻ 126.9 mM; Ca²⁺ 1.6 mM; Mg²⁺ 1.2 mM; H₂PO₄⁻ 1.2 mM; SO₄²⁻ 1.2 mM; HCO₃⁻ 25.0 mM; D-glucose 11.1 mM) of pH 7.4. Additionally, 1 µmol/l of indomethacin were added to the Krebs-Henseleit solution to inhibit prostaglandin biosynthesis. The rings were subsequently precontracted by addition of phenylephrine (concentration in the solution: 1 µM) and the endothelium-dependent relaxation or the functional loss of the endothelium was tested by addition of 20 acetylcholine (concentration in the solution: 1 µM). After a 30-minute washing period, the rings were then again precontracted by addition of phenylephrine (1 µM), and the relaxing action of the test substances of the formula I was determined by administration of cumulative doses of the latter. The data were evaluated by standard methods. Given is the concentration IC₅₀ by which contraction is inhibited by 50 % (50 % relaxation).

25

The following results were obtained:

Compound of

IC₅₀

Example No.

	59	ring without endothelium	0.27 μ M
	59	ring with endothelium	0.52 μ M
5	88	ring without endothelium	0.29 μ M
	88	ring with endothelium	0.67 μ M
	129	ring without endothelium	0.31 μ M
	129	ring with endothelium	0.46 μ M

(3) Hemodynamic effect in the pig

Three pigs (German landrace) were anesthetized (Ketamine 20 mg/kg i.m., Methomidate 8 mg/kg i.p., Xylazine 2.5 mg/kg i.m. and Pentobarbital 25 mg/kg i.v. as a bolus plus 0.16 mg/kg per minute). The trachea was intubated and the animals were given artificial respiration with air. Oxygen was added to keep the blood gas parameters in the normal range. To record the blood pressure (BP; BP(s) = systolic blood pressure, BP(d) = diastolic blood pressure) by means of a Statham 23Db pressure transducer a catheter was inserted into the right A. Femoralis. The left ventricular pressure (LVP), the left ventricular end-diastolic pressure (LVEDP), the contractility (dP/dt) and the heart rate (HR) were determined with a Millar PC 350 catheter "tip manometer" which was inserted into the right ventricle. After a stabilization period of the hemodynamic parameters of 30 minutes the test substance was administered in the indicated dose into the exposed duodenum by means of a catheter. The determined data were evaluated according to standard methods. Given are the means and the standard deviations ($M \pm SEM$) of the starting values and of the maximal changes of the individual parameters (= maximal effects).

The following results were obtained:

Compound of example 88 (dosage 10 mg/kg i.d.)

5	Parameter	starting value	maximal alteration	duration of action (min)
	BP(s) (mm Hg)	123 ± 26	-23 ± 6	> 180
	BP(d) (mm Hg)	83 ± 24	-20 ± 8	> 180
10	LVEDP (mm Hg)	4 ± 0.6	-1.3 ± 0.3	> 180
	dP/dt _{max} (mm Hg/sec))	1800 ± 289	-633 ± 33	> 180
	HR (beats/min)	98 ± 2	-8 ± 2	> 180

The invention may be embodied in other specific forms and those skilled in the art will recognize that various changes and modifications can be made without departing from the spirit or essential characteristics thereof. The present embodiment is, therefore, considered in all respects as illustrative and not restrictive, the scope of the appended claims rather than by the foregoing description, and all changes which come within the meaning and range of equivalency of the claims are therefore intended to be embraced therein.